

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 634 169 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
05.01.2000 Bulletin 2000/01

(51) Int. Cl.⁷: **A61K 31/395**, A61K 31/495,
A61K 31/47, A61K 31/435,
A61K 31/505

(21) Application number: **94109861.8**

(22) Date of filing: **25.06.1994**

(54) **Quinoline or quinazoline derivatives and their use in the manufacture of a medicament for the treatment of osteoporosis**

Chinoline oder Chinazolin-Derivate und deren Verwendung zur Herstellung eines Medikaments für die Behandlung von Osteoporose

Dérivés quinolines ou quinazolines et leur utilisation dans la fabrication d'un médicament pour le traitement de l'ostéoporose

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT SE

(30) Priority: **29.06.1993 JP 15865293**

(43) Date of publication of application:
18.01.1995 Bulletin 1995/03

(73) Proprietor:
TAKEDA CHEMICAL INDUSTRIES, LTD.
Chuo-ku, Osaka 541 (JP)

(72) Inventors:
• **Sohda, Takashi**
Osaka 569 (JP)

• **Taketomi, Shigehisa**
Ikeda, Osaka 563 (JP)
• **Baba, Atsuo**
Ashiya, Hyogo 659 (JP)

(74) Representative:
von Kreisler, Alek, Dipl.-Chem. et al
Patentanwälte,
von Kreisler-Selting-Werner,
Bahnhofsvorplatz 1 (Deichmannhaus)
50667 Köln (DE)

(56) References cited:
EP-A- 0 403 185 **EP-A- 0 567 107**
EP-A- 0 608 870 **US-A- 4 343 940**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 634 169 B1

Description

[0001] The present invention relates to a pharmaceutical composition containing a quinoline or quinazoline derivative which has inhibitory activity of bone resorption and is useful as a prophylactic or therapeutic agent against osteoporosis.

BACKGROUND OF THE INVENTION

[0002] Osteoporosis is a morbid state or disease wherein bone mass is so decreased as to cause some symptoms or danger. Its main symptoms are kyphosis of spine and fracture of lumbar vertebrae, thoracic vertebrae, femoral neck, distal ends of radii, ribs, the proximal ends of humeri or the like. In normal bone tissues, destruction by bone formation and bone resorption are repeated with a balance. Osteoblasts and osteoclasts play major roles in bone formation and bone resorption, respectively. When the balance between bone formation and bone resorption is lost and bone resorption exceeds bone formation, bone mass is decreased. Therefore, it is believed that drugs inhibiting bone resorption are useful for preventing and treating osteoporosis, and bone resorption inhibitors such as estrogen, calcitonin and the like have been administered as drugs for treating osteoporosis.

[0003] However, in the administration of these drugs, the subject is limited and the resulting effects are sometimes uncertain, and satisfactory effects have not been obtained.

[0004] Thus, it is desired to develop an agent of preventing or treating the increase of bone resorption.

[0005] EP-A-0 567 107 describes quinoline and quinazoline derivatives for treating arthritis.

OBJECTS OF THE INVENTION

[0006] The main object of the present invention is to provide a pharmaceutical composition for inhibiting bone resorption.


[0007] Another object of the present invention is to provide a pharmaceutical composition for preventing or treating osteoporosis.

[0008] These objects as well as other objects and advantages of the present invention will become apparent to those skilled in the art from the following description.

SUMMARY OF THE INVENTION

[0009] The present inventors have intensively studied to develop generally applicable pharmaceutical compositions which have a direct effect on bones to inhibit bone resorption. As a result, it has been found that quinoline or quinazoline derivatives of the following formula (I) have a direct effect on bones to exhibit potent inhibitory activity of bone resorption.

[0010] Thus, the present invention is directed to the use of a compound of the formula (I):



wherein

Y is a nitrogen atom or C-G in which G is an optionally esterified or optionally amidated carboxyl group, or hydroxymethyl group;

R is an optionally substituted hydrocarbon group or optionally substituted heterocyclic group;

X is an oxygen atom or optionally oxidized sulfur atom;

n is 0 or 1;

k is 0 or 1;


G and R may be linked together to form a ring;

each of the ring A and ring B may optionally be substituted;

or a pharmaceutically acceptable salt thereof for formulation of a pharmaceutical composition for inhibiting bone resorption.

[0011] The present invention is also directed to the use of a compound of the above formula (I) or a pharmaceutically acceptable salt thereof, for formulation of a pharmaceutical composition for preventing or treating osteoporosis.

[0012] The present invention is furthermore directed to a compound of the formula (I'):



wherein

Y is C-G in which G is an optionally amidated carboxyl group, or hydroxymethyl group;

R is an optionally substituted hydrocarbon group or optionally substituted heterocyclic group;

X is an oxygen atom or optionally oxidized sulfur atom;

n is 0 or 1;

k is 0 or 1;

G and R may be linked together to form a ring;


each of the ring A and B may optionally be substituted;

or a pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION OF THE INVENTION

[0013] Several compounds included in the compound (I) used in the present invention are novel and the present applicant have already filed on these compounds (Japanese Patent Application Nos. 5-012628 and 5-095780).

[0014] The optionally substituted hydrocarbon residue represented by R is preferably a group of the formula:



wherein X^1 is an oxygen atom, optionally oxidized sulfur atom or $-(\text{CH}_2)_m-$ (wherein m is an integer of 0 to 5), and Z^1 is an optionally substituted hydrocarbon group, optionally substituted heterocyclic group or optionally substituted amino group.

[0015] Examples of the hydrocarbon group of the optionally substituted hydrocarbon group represented by R or Z^1 in the above formula (I) include aliphatic hydrocarbon groups, alicyclic hydrocarbon groups, alicyclic-aliphatic hydrocarbon groups, (aromatic carbocycle)-aliphatic hydrocarbon groups, aromatic hydrocarbon groups and the like.

[0016] Examples of such aliphatic hydrocarbon groups include saturated aliphatic hydrocarbon groups having 1 to 8 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, t-butyl, pentyl, isopentyl, neopentyl, t-pentyl, hexyl, isohexyl, heptyl, octyl and the like; unsaturated aliphatic hydrocarbon groups having 2 to 8 carbon atoms such as ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-methyl-1-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 3-methyl-2-butenyl, 1-hexenyl, 3-hexenyl, 2,4-hexadienyl, 5-hexenyl, 1-heptenyl, 1-octenyl, ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-hexynyl, 3-hexynyl, 2,4-hexadiynyl, 5-hexynyl, 1-heptynyl, 1-octynyl and the like.

[0017] Examples of such alicyclic hydrocarbon groups include saturated alicyclic hydrocarbon groups having 3 to 7 carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like; and unsaturated alicyclic hydrocarbon groups having 5 to 7 carbon atoms such as 1-cyclopentenyl, 2-cyclopentenyl, 3-cyclopentenyl, 1-cyclohexenyl, 2-cyclohexenyl, 3-cyclohexenyl, 1-cycloheptenyl, 2-cycloheptenyl, 3-cycloheptenyl, 2,4-cycloheptadienyl

and the like.

[0018] Examples of such alicyclic-aliphatic hydrocarbon groups include those having 4 to 9 carbon atoms each of which is composed of the above alicyclic hydrocarbon group and aliphatic hydrocarbon group, for example, cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl, 2-cyclopentenylmethyl, 3-cyclopentenylmethyl, cyclohexylmethyl, 2-cyclohexenylmethyl, 3-cyclohexenylmethyl, cyclohexylethyl, cyclohexylpropyl, cycloheptylmethyl, cycloheptylethyl and the like.

[0019] Examples of such (aromatic carbocycle)-aliphatic hydrocarbon groups include phenyl alkyl having 7 to 9 carbon atoms such as benzyl, phenethyl, 1-phenylethyl, 3-phenylpropyl, 2-phenylpropyl, 1-phenylpropyl and the like; naphthylalkyl having 11 to 13 carbon atoms such as α -naphthylmethyl, α -naphthylethyl, β -naphthylmethyl, β -naphthylethyl and the like.

[0020] Examples of such aromatic hydrocarbon groups include phenyl, naphthyl (e.g., α -naphthyl, β -naphthyl) and the like.

[0021] Examples of the oxidized sulfur atom represented by X^1 include a thio group, sulfinyl group and sulfonyl group. In particular, a thio group is preferred.

[0022] X^1 is preferably $-(CH_2)_m-$ wherein m is 1 or 2.

[0023] Examples of the optionally substituted hydrocarbon group represented by Z^1 include the same optionally substituted hydrocarbon groups as those described above for R.

[0024] Examples of the optionally substituted heterocyclic group represented by Z^1 include the same optionally substituted heterocyclic groups as those described below for R. It is preferably aromatic 5-membered heterocyclic groups containing 2 or 3 heteroatoms (e.g., oxygen atom, nitrogen atom and sulfur atom), more preferably 1,2,4-triazol-1-yl.

[0025] Examples of the heterocyclic group of the optionally substituted heterocyclic group represented by R or Z^1 in the above formula (I) include 5- to 7-membered heterocyclic groups containing one sulfur atom, nitrogen atom or oxygen atom; 5- to 6-membered heterocyclic groups containing 2 to 4 nitrogen atoms; 5- to 6-membered heterocyclic groups containing 1 to 2 nitrogen atoms, and one sulfur atom or oxygen atom. Each of these heterocyclic group may form a condensed ring with a 6-membered ring containing up to 2 nitrogen atoms, benzene ring or 5-membered ring containing one sulfur atom.

[0026] Specific examples of the heterocyclic group include 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrazinyl, 2-pyrrolyl, 3-pyrrolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-pyrazolyl, 4-pyrazolyl, isothiazolyl, isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 1,2,4-triazol-3-yl, 1,3,4-triazol-2-yl, 1,2,3-triazol-4-yl, tetrazol-5-yl, benzimidazol-2-yl, indol-3-yl, benzpyrazol-3-yl, 1H-pyrrolo[2,3-b]pyrazin-2-yl, 1H-pyrrolo[2,3-b]pyridin-6-yl, 1H-imidazo[4,5-b]pyridin-2-yl, 1H-imidazo[4,5-c]pyridin-2-yl, 1H-imidazo[4,5-b]pyrazin-2-yl and the like.

[0027] Each of the hydrocarbon groups and heterocyclic groups represented by R or Z^1 in the above formula (I) may be unsubstituted or have 1 to 3 substituents at any possible position in the ring. Such substituents include, for example, aliphatic chain hydrocarbon groups, alicyclic hydrocarbon groups, aryl groups, aromatic heterocyclic groups, non-aromatic heterocyclic groups, halogen atoms, nitro group, optionally substituted amino group, optionally substituted acyl groups, optionally substituted hydroxyl group, optionally substituted thiol group, optionally esterified carboxyl group and the like.

[0028] The aliphatic chain hydrocarbon group as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z^1 include, for example, straight chain or branched chain aliphatic hydrocarbon groups such as alkyl groups, preferably alkyl groups having 1 to 10 carbon atoms; alkenyl groups, preferably alkenyl groups having 2 to 10 carbon atoms; alkynyl groups and the like.

[0029] Preferred examples of the alkyl group include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, tert-pentyl, 1-ethylpropyl, hexyl, isohexyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 2-ethylbutyl, hexyl, pentyl, octyl, nonyl, decyl and the like.

[0030] Preferred examples of the alkenyl include vinyl, allyl, isopropenyl, 1-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 2-ethyl-1-butenyl, 3-methyl-2-butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 4-methyl-3-pentenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl and the like.

[0031] Preferred examples of the alkynyl include ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-pentylnyl, 4-pentylnyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl and the like.

[0032] The alicyclic hydrocarbon group as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z^1 include, for example, saturated or unsaturated alicyclic hydrocarbon groups such as cycloalkyl, cycloalkenyl, cycloalkadienyl and the like.

[0033] Preferred examples of the cycloalkyl include that having 3 to 10 carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl, bicyclo[3.2.1]octyl, bicyclo[3.2.2]nonyl, bicyclo[3.3.1]nonyl, bicyclo[4.2.1]nonyl, bicyclo[4.3.1]decyl and the like.

[0034] Preferred examples of the cycloalkenyl include that having 5 to 7 carbon atoms such as 2-cyclopenten-1-yl, 3-cyclopenten-1-yl, 2-cyclohexen-1-yl, 3-cyclohexen-1-yl and the like.

[0035] Preferred examples of the cycloalkadienyl include that having 5 to 7 carbon atoms such as 2,4-cyclopentadien-1-yl, 2,4-cyclohexadien-1-yl, 2,5-cyclohexadien-1-yl and the like.

[0036] The aryl group as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z¹ means a monocyclic or condensed polycyclic aromatic hydrocarbon group. Preferred examples thereof include phenyl, naphthyl, anthryl, phenanthryl, acenaphthylenyl and the like. More preferred examples thereof are phenyl, 1-naphthyl, 2-naphthyl and the like.

[0037] Preferred examples of the aromatic heterocyclic group as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z¹ include aromatic monocyclic heterocyclic groups such as furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, furazanyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, tetrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl and the like; aromatic condensed heterocyclic groups such as benzofuranyl, isobenzofuranyl, benzo[b]thienyl, indolyl, isoindolyl, 1H-indazolyl, benzimidazolyl, benzoxazolyl, 1,2-benzisoxazolyl, benzothiazolyl, 1,2-benzisothiazolyl, 1H-benzotriazolyl, quinolyl, isoquinolyl, cinnolyl, quinazolyl, quinoxalyl, phthalazinyl, naphthyridinyl, purinyl, pteridinyl, carbazolyl, α -carbolinyl, β -carbolinyl, γ -carbolinyl, acridinyl, phenoxazinyl, phenothiazinyl, phenazinyl, phenoxathiinyl, thianthrenyl, phenanthridinyl, phenanthrolinyl, indolizinyl, pyrrolo[1,2-b]pyridazinyl, pyrazolo[1,5-a]pyridyl, imidazo[1,2-a]pyridyl, imidazo[1,5-a]pyridyl, imidazo[1,2-b]pyridazinyl, imidazo[1,2-a]pyrimidinyl, 1,2,4-triazolo[4,3-a]pyridyl, 1,2,4-triazolo[4,3-b]pyridazinyl and the like.

[0038] Preferred examples of the non-aromatic heterocyclic group as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z¹ include oxiranyl, azetidiny, oxetanyl, thietanyl, pyrrolidinyl, tetrahydrofuranyl, thiolanyl, piperidinyl, tetrahydropyranyl, morpholyl, thiomorpholyl, piperazinyl and the like.

[0039] The halogen as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z¹ includes, for example, fluorine, chlorine, bromine and iodine. In particular, fluorine and chlorine are preferred.

[0040] The optionally substituted amino group as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z¹ includes, for example, an amino group and an amino group substituted with one or two substituents such as alkyl having 1 to 10 carbon atoms, alkenyl having 1 to 10 carbon atoms or aromatic groups (e.g., methylamino, dimethylamino, ethylamino, diethylamino, dibutylamino, diallylamino, cyclohexylamino, phenylamino, N-methyl-N-phenylamino, etc.).

[0041] The optionally substituted acyl group as the substituent of hydrocarbon group or heterocyclic group represented by R or Z¹ includes, for example, formyl, and (C₁₋₁₀ alkyl)-carbonyl, (C₁₋₁₀ alkenyl)-carbonyl and aromatic carbonyl (e.g., acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, heptanoyl, octanoyl, cyclobutanecarbonyl, cyclopentanecarbonyl, cyclohexanecarbonyl, cycloheptanecarbonyl, crotonyl, 2-cyclohexenecarbonyl, benzoyl, nicotinoyl, etc.).

[0042] The optionally substituted hydroxyl group of the substituent of the hydrocarbon group or heterocyclic group represented by R or Z¹ includes, for example, a hydroxyl group and a hydroxyl group having an appropriate substituent such as a protecting group for a hydroxyl group. Examples of the hydroxyl group having such a substituent include alkoxy, alkenyloxy, aralkyloxy, acyloxy, aryloxy and the like.

[0043] Preferred examples of the alkoxy include that having 1 to 10 carbon atoms such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy, isopentyloxy, neopentyloxy, hexyloxy, heptyloxy, nonyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy and the like.

[0044] Examples of the alkenyloxy include that having 1 to 10 carbon atoms such as allyloxy, crotyloxy, 2-pentenyl, 3-hexenyloxy, 2-cyclopentenylmethoxy, 2-cyclohexenylmethoxy and the like.

[0045] Examples of the aralkyloxy include phenyl-C₁₋₄ alkyloxy (e.g., benzyloxy, phenethyloxy, etc.).

[0046] Preferred examples of the acyloxy include alkanoyloxy having 2 to 4 carbon atoms such as acetyloxy, propionyloxy, n-butyryloxy, iso-butyryloxy and the like.

[0047] Examples of the aryloxy include phenoxy, 4-chlorophenoxy and the like.

[0048] The optionally substituted thiol group as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z¹ includes, for example, a thiol group and a thiol group having an appropriate substituent such as a protecting group for a thiol group. Examples of the thiol group having such a substituent include alkylthio, aralkylthio, acylthio and the like.

[0049] Preferred examples of the alkylthio include alkylthio having 1 to 10 carbon atoms such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio, sec-butylthio, tert-butylthio, pentylthio, isopentylthio, neopentylthio, hexylthio, heptylthio, nonylthio, cyclobutylthio, cyclopentylthio, cyclohexylthio and the like.

[0050] Examples of the aralkylthio include phenyl-C₁₋₄ alkylthio (e.g., benzylthio, phenethylthio, etc.).

[0051] Preferred examples of the acylthio include alkanoylthio having 2 to 4 carbon atoms such as acetylthio, propionylthio, n-butyrylthio, iso-butyrylthio and the like.

[0052] The optionally esterified carboxyl group as the substituent of the hydrocarbon group or heterocyclic group represented by R or Z¹ include, for example, a carboxyl group, (C₁₋₆ alkoxy)-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-

butoxycarbonyl, pentyloxycarbonyl, hexyloxycarbonyl, etc.), (C₃₋₆ alkenyloxy)-carbonyl (e.g., allyloxycarbonyl, crotyloxycarbonyl, 2-pentenylloxycarbonyl, 3-hexenylloxycarbonyl, etc.), aralkyloxycarbonyl (e.g., benzyloxycarbonyl, phenethylloxycarbonyl, etc.) and the like.

[0053] The substituent of the hydrocarbon group and heterocyclic group represented by R or Z¹ in the above formula (I) may have at least one, preferably 1 to 3 appropriate substituents. Such substituents include, for example, lower alkyl groups, lower alkenyl groups, lower alkynyl groups, cycloalkyl groups, aryl groups, aromatic heterocyclic groups, non-aromatic heterocyclic groups, aralkyl groups, an amino group, N-monosubstituted amino groups, N,N-disubstituted amino groups, an amidino group, acyl groups, a carbamoyl group, N-monosubstituted carbamoyl groups, N,N-disubstituted carbamoyl groups, a sulfamoyl group, N-monosubstituted sulfamoyl groups, N,N-disubstituted sulfamoyl groups, a carboxyl group, lower alkoxycarbonyl, a hydroxyl group, lower alkoxy groups, lower alkenyloxy groups, cycloalkyloxy groups, lower alkylthio groups, aralkylthio groups, arylthio groups, a sulfo group, a cyano group, an azido group, halogen atoms, a nitro group, a nitroso group and the like. Specific examples of these substituents include those described above with respect to the substituents of the hydrocarbon group and heterocyclic group represented by R or Z¹.

[0054] When R is -CH₂-X¹-Z¹ in the formula (I), the optionally substituted amino group represented by Z¹ is represented by the formula: -N(R¹)(R²) wherein R¹ and R² are the same or different and are hydrogen, optionally substituted hydrocarbon group or optionally substituted heterocyclic group.

[0055] The hydrocarbon group in the optionally substituted hydrocarbon group and heterocyclic group in the optionally substituted heterocyclic group represented by R¹ or R² include, for example, the same hydrocarbon groups and heterocyclic groups as those represented by R.

[0056] R¹ and R² may be linked together to form a ring. Examples of such -N(R¹)(R²) include 1-pyrrolidinyl, 1-imidazolidinyl, 1-pyrazolidinyl, 1-piperidinyl, 1-piperazinyl, 4-morpholinyl, 4-thiomorpholinyl, homopiperazin-1-yl, 1,2,4-triazol-1-yl, 1,3,4-triazol-1-yl, pyrazol-1-yl, imidazol-1-yl, 1,2,3-triazol-1-yl, 1,2,3-triazol-2-yl, tetrazol-1-yl, benzimidazol-1-yl, indol-1-yl, 1H-indazol-1-yl and the like.

[0057] Each of the hydrocarbon group and heterocyclic group represented by R¹ or R² may have 1 to 3 substituents at any possible position in the ring. Examples of the substituent include the same substituents as those of the hydrocarbon group or heterocyclic group represented by R. These substituents of the hydrocarbon group and heterocyclic group represented by R¹ or R² each may have at least one, preferably 1 to 3 appropriate substituents. Such substituents include, for example, lower alkyl groups, lower alkenyl groups, lower alkynyl groups, cycloalkyl groups, aryl groups, aromatic heterocyclic groups, non-aromatic heterocyclic groups, aralkyl groups, an amino group, N-monosubstituted amino groups, N,N-disubstituted amino groups, an amidino group, acyl groups, a carbamoyl group, N-monosubstituted carbamoyl groups, N,N-disubstituted carbamoyl groups, a sulfamoyl group, N-monosubstituted sulfamoyl groups, N,N-disubstituted sulfamoyl groups, a carboxyl group, lower alkoxycarbonyl groups, a hydroxyl group, lower alkoxy groups, lower alkenyloxy groups, cycloalkyloxy groups, aralkyloxy groups, aryloxy groups, a mercapto group, lower alkylthio groups, aralkylthio groups, arylthio groups, a sulfo group, a cyano group, an azido group, a nitro group, a nitroso group, halogen and the like. Specific examples of these substituents include those described above with respect to the substituents of the hydrocarbon group and heterocyclic group represented by R.

[0058] Examples of the optionally oxidized sulfur atom represented by X include a thio group, sulfinyl group and sulfonyl group. In particular, a thio group is preferred.

[0059] Each of the ring A and ring B in the formula (I) may be substituted with at least one substituent. Examples of the substituent include halogen atoms, a nitro group, optionally substituted alkyl groups, an optionally substituted hydroxyl group, an optionally substituted thiol group, an optionally substituted amino group, optionally substituted acyl groups, an optionally esterified carboxyl group and optionally substituted aromatic cyclic groups.

[0060] The halogen atom as the substituent of the ring A and ring B includes, for example, fluorine, chlorine, bromine and iodine. In particular, fluorine and chlorine are preferred.

[0061] The optionally substituted alkyl group as the substituent of the ring A and ring B may be, for example, any of straight-chain, branched-chain or cyclic alkyl groups. Specific examples thereof include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl, nonyl, decyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like.

[0062] The optionally substituted hydroxyl group as the substituent of the ring A and ring B includes, for example, a hydroxyl group and a hydroxyl group having an appropriate substituent such as that used as a protecting group for a hydroxyl group (e.g., alkoxy, alkenyloxy, aralkyloxy, acyloxy, aryloxy, etc.).

[0063] Preferred examples of the alkoxy include that having 1 to 10 carbon atoms such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy, isopentyloxy, neopentyloxy, hexyloxy, heptyloxy, nonyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy and the like.

[0064] Examples of the alkenyloxy include that having 1 to 10 carbon atoms such as allyloxy, crotyloxy, 2-pentenylloxy, 3-hexenylloxy, 2-cyclopentenylmethoxy, 2-cyclohexenylmethoxy and the like.

[0065] Examples of the aralkyloxy include phenyl-C₁₋₄ alkylloxy (e.g., benzyloxy, phenethyloxy, etc.).

[0066] Preferred examples of the acyloxy include alkanoyloxy having 2 to 4 carbon atoms such as acetyloxy, propio-

nyloxy, n-butyryloxy, iso-butyryloxy and the like.

[0067] Examples of the aryloxy include phenoxy, 4-chlorophenoxy and the like.

[0068] The optionally substituted thiol as the substituent of the ring A and ring B includes, for example, a thiol group and a thiol group having an appropriate substituent such as that used as a protecting group for a thiol group (e.g., alkylthio, aralkylthio, acylthio, etc.).

[0069] Preferred examples of the alkylthio include that having 1 to 10 carbon atoms such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio, sec-butylthio, tert-butylthio, pentylthio, isopentylthio, neopentylthio, hexylthio, heptylthio, nonylthio, cyclobutylthio, cyclopentylthio, cyclohexylthio and the like.

[0070] Examples of the aralkylthio include phenyl-C₁₋₄ alkylthio (e.g., benzylthio, phenethylthio, etc.).

[0071] Preferred examples of the acylthio include alkanoylthio having 2 to 4 carbon atoms such as acetylthio, propionylthio, n-butyrylthio, iso-butyrylthio and the like.

[0072] The optionally substituted amino group as the substituent of the ring A and ring B includes, for example, an amino group and an amino group substituted with one or two substituents selected from alkyl groups having 1 to 10 carbon atoms, alkenyl groups having 1 to 10 carbon atoms and aromatic groups. Specific examples of the substituted amino group include methylamino, dimethylamino, ethylamino, diethylamino, dibutylamino, diallylamino, cyclohexylamino, phenylamino, N-methyl-N-phenylamino and the like.

[0073] The acyl group as the substituent of the ring A and ring B include, for example, formyl, and (C₁₋₁₀ alkyl)-carbonyl, (C₁₋₁₀ alkenyl)-carbonyl and (aromatic group)-carbonyl such as acetyl, propionyl, butyryl, isobutyryl, valeryl, pivaloyl, hexanoyl, heptanoyl, octanoyl, cyclobutanecarbonyl, cyclopentanecarbonyl, cyclohexanecarbonyl, cycloheptanecarbonyl, crotonyl, 2-cyclohexenecarbonyl benzoyl, nicotinoyl and the like.

[0074] The optionally esterified carboxyl group as the substituent of the ring A and ring B includes, for example, a carboxyl group, (C₁₋₆ alkyloxy)-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl, hexyloxycarbonyl, etc.), (C₃₋₆ alkenyloxy)-carbonyl (e.g., allyloxycarbonyl, crotyloxycarbonyl, 2-pentenylloxycarbonyl, 3-hexenyloxycarbonyl, etc.), aralkyloxycarbonyl (e.g., benzyloxycarbonyl, phenethylloxycarbonyl, etc.) and the like.

[0075] The aromatic cyclic group as the substituent of the ring A and ring B includes, for example, C₆₋₁₄ aromatic hydrocarbon groups (e.g., phenyl, naphthyl, anthryl, etc.) and aromatic heterocyclic groups (e.g., pyridyl, furyl, thienyl, imidazolyl, thiazolyl, etc.).

[0076] The substituent of the ring A and ring B may be at any possible position in each ring. The each ring may be substituted with the same or different 1 to 4 substituents. When the substituents of the ring A or ring B are adjacent to each other, the adjacent substituents are linked together to form a group of the formula: -(CH₂)_t- or -O-(CH₂)₁-O- (wherein t is an integer of 3 to 5 and 1 is an integer of 1 to 3) which may form a 5- to 7-membered ring with carbon atoms of the benzene ring.

[0077] Preferably, the ring A is substituted with at least one alkoxy group, preferably at least one methoxy group; or the same or different two alkoxy groups, preferably two methoxy groups. More preferably, the ring A is substituted with two methoxy groups at the 6- and 7-positions of the quinoline ring or quinazoline ring.

[0078] Preferably, the ring B is substituted with at least one alkoxy group, preferably at least one methoxy group or isopropoxy group; or the same or different two alkoxy groups. More preferably, the ring B is substituted with an isopropoxy group at the 3-position and a methoxy group at the 4-position.



[0079] When the compound of the formula (I) is a quinoline derivative wherein Y is C-G, the optionally esterified carboxyl group represented by G may be, for example, a carboxyl group, alkyloxycarbonyl groups, aralkyloxycarbonyl groups and the like.

[0080] The alkyl group of the alkyloxycarbonyl group includes, for example, alkyl groups having 1 to 6 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl and the like.

[0081] The aralkyl group of the aralkyloxycarbonyl group means an alkyl group having an aryl group as a substituent (i.e., arylalkyl group). Examples of the aryl group include phenyl, naphthyl and the like. The aryl group may have the same substituent as that of the above ring A. The alkyl group is preferably a lower alkyl group having 1 to 6 carbon atoms. Preferred examples of the aralkyl group include benzyl, phenethyl, 3-phenylpropyl, (1-naphthyl)methyl, (2-naphthyl)methyl and the like. In particular, benzyl, phenethyl and the like are preferred.

[0082] When G is an amidated carboxyl group, the amidated carboxyl group is represented by -CON(R¹)(R²) wherein each symbol is as defined above.

[0083] When the compound of the formula (I) is a quinoline derivative wherein Y is C-G, R and G may be linked together to form a 5-membered ring. The resulting compound is represented, for example, by the following formulas (II) or (III).



wherein R^3 is hydrogen, an optionally substituted hydrocarbon group or optionally substituted heterocyclic group, and the other symbols are as defined above.

[0084] The optionally substituted hydrocarbon group and optionally substituted heterocyclic group represented by R^3 in the formulas (II) and (III) includes, for example, the same groups as described for R or Z^1 .

[0085] Y in the formula (I) is preferably C-G. More preferably, G is a (C_{1-6} alkyl)oxycarbonyl group, particularly preferably an ethoxycarbonyl group.

[0086] n in the formula (I) is preferably 0.

[0087] k in the formula (I) is preferably 0.

[0088] Preferred examples of the compound of the formula (I) include:

methyl 4-(3,4-dimethoxyphenyl)-2-ethyl-6,7-dimethoxyquinoline-3-carboxylate;
ethyl 6-chloro-2-methyl-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate;
6,7-dimethoxy-9-phenylfuro[3,4-b]quinoline-1(3H)-one;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate;
4-(3,4-dimethoxyphenyl)-2-(2-hydroxyethylthiomethyl)-6,7-dimethoxyquinazoline;
4-(3,4-diethoxyphenyl)-6,7-dimethoxy-2-[(4-methyl-1,2,4-triazol-3-yl)thiomethyl]quinazoline;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[2-(1-methylimidazol-2-yl)ethyl]quinoline-3-carboxylate;
methyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-3-methoxycarbonylquinoline-2-acetate;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
ethyl 4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
ethyl 4-(4-hydroxy-3-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
ethyl 7-hydroxy-6-methoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate 1-oxide; and
ethyl 2-(N,N-diethylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate.

[0089] In particular, the compound of the formula (I) is preferably that wherein Y is C-G in which G is ethoxycarbonyl, R is $-CH_2-Z^1$, Z^1 is 1,2,4-triazol-1-yl, the ring A is substituted with methoxy groups at the 6- and 7-positions of the quinoline ring, the ring B is substituted with methoxy groups at the 3- and 4-positions, n is 0, and k is 0.

[0090] The salt of the compound of the formula (I) is preferably a pharmaceutically acceptable salt. Examples thereof include salts with inorganic bases, organic bases, inorganic acids, organic acids, basic or acidic amino acids and the like.

[0091] Preferred examples of the salts with inorganic bases include alkaline metal salts such as a sodium salt, potassium salt and the like; alkaline earth metal salts such as a calcium salt, magnesium salt and the like; an aluminium salt; an ammonium salt and the like.

[0092] Preferred examples of the salts with organic bases include salts with trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, N,N'-dibenzylethylenediamine and the like.

[0093] Preferred examples of the salts with inorganic acids include salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid and the like.

[0094] Preferred examples of the salts with organic acids include salts with formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid and the like.

[0095] Preferred examples of the salts with basic amino acids include salts with arginine, lysine, ornithine and the like. Preferred examples of the salts with acidic amino acids include aspartic acid, glutamic acid and the like.

[0096] The compound of the formula (I) can be formulated with a pharmaceutically acceptable carrier and administered orally or parenterally as solid preparations such as tablets, capsules, granules, powders or the like; or liquid preparations such as syrups, injections or the like.

[0097] As the pharmaceutically acceptable carrier, various organic or inorganic carrier materials conventionally used for pharmaceutical preparations can be used and formulated as excipients, lubricants, binders, disintegrators and the like for solid preparations; solvents, solution adjuvants, suspending agents, tonicity agents, buffering agents, soothing agents and the like for liquid preparations and the like. If necessary, pharmaceutical additives such as antiseptics, antioxidants, colorants, sweetening agents and the like can be used.

[0098] Preferred examples of the excipient include lactose, sucrose, D-mannitol, starch, crystalline cellulose, light anhydrous silicic acid and the like.

[0099] Preferred examples of the lubricant include magnesium stearate, calcium stearate, talc, colloidal silica and the like.

[0100] Preferred examples of the binder include crystalline cellulose, sucrose, D-mannitol, dextrin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, polyvinyl-pyrrolidone and the like.

[0101] Preferred examples of the disintegrator include starch, carboxymethylcellulose, carboxymethyl cellulose calcium, croscarmellose sodium, carboxymethyl starch sodium and the like.

[0102] Preferred examples of the solvent include water for injection, alcohols, propylene glycol, macrogol, sesame oil, corn oil and the like.

[0103] Preferred examples of the solution adjuvant include polyethylene glycol, propylene glycol, D-mannitol, benzyl benzoate, ethanol, trisaminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate and the like.

[0104] Preferred examples of the suspending agent include surfactants such as stearyl triethanolamine, sodium lauryl sulfate, laurylaminopropionic acid, lecithin, benzalkonium chloride, benzethonium chloride, glyceryl monostearate and the like; hydrophilic polymers such as polyvinyl alcohol, polyvinyl pyrrolidone, carboxymethyl cellulose sodium, methyl cellulose, hydroxymethyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose and the like.

[0105] Preferred examples of the tonicity agent include sodium chloride, glycerin, D-mannitol and the like.

[0106] Preferred examples of the buffering agent include buffers such as phosphates, acetates, carbonates, citrates and the like.

[0107] Preferred examples of the soothing agent include benzyl alcohol and the like.

[0108] Preferred examples of the antiseptics include parahydroxybenzoic acid esters, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, sorbic acid and the like.

[0109] Preferred examples of the antioxidant include sulfites, ascorbic acid and the like.


[0110] The compound (I) used in the present invention has low toxicity. For example, the compounds of Examples 142 and 181 are orally administered to mice in a dose of 1,000 mg/kg, no mouse was died. Thus, the compound (I) is a useful agent as a bone resorption inhibitor for preventing or treating osteoporosis for mammals such as humans, cattle, horses, swine, dogs, cats and the like.

[0111] The dose of the compound (I) used in the present invention can be appropriately selected depending upon the administration route and condition of the patient to be treated. Normally, the dose can be selected from the regions of 10 mg to 500 mg per adult in the case of oral administration and 1 mg to 100 mg per adult in the case of parenteral administration. The compound in the above dose can be administered a day in one to three divided doses.

[0112] The above compound (I) can be prepared, for example, as follows.

Method A

[0113]



wherein Q is an optionally substituted hydrocarbon group, R^4 is a lower alkyl group, and the other symbols are as defined above.

[0114] The optionally substituted hydrocarbon group by Q in the formulas (I-1) and (V-1) includes the same groups as those described above for R or Z^1 .


[0115] Examples of the lower alkyl group represented by R^4 in the formulas (I-1) and (V-1) include alkyl groups having 1 to 4 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl and the like.

[0116] In this method, a 2-aminobenzophenone derivative (IV) is reacted with the compound (V-1) in the presence of an acid to prepare the compound (I-1). The reaction of the compound (IV) and the compound (V-1) can be carried out in an appropriate solvent. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), ethers (e.g., dioxane, tetrahydrofuran, dimethoxyethane, etc.), alcohols (e.g., methanol, ethanol, propanol, etc.), N,N-dimethylformamide, dimethyl sulfoxide, chloroform, dichloromethane, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane, acetic acid and the like. The reaction of the compound (IV) and the compound (V-1) is carried out in the presence of an appropriate acid such as Lewis acids (e.g., aluminium chloride, zinc chloride, etc.), hydrochloric acid, sulfuric acid, trifluoroacetic acid, p-toluenesulfonic acid or the like. The amount of the acid to be used is preferably 0.05 to 2.0 mol per mol of the compound (IV). The reaction temperature is normally 20°C to 200°C, preferably about 30°C to 150°C. The reaction time is 0.5 to 20 hours, preferably 1 to 10 hours.

[0117] The compound (I-1) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method B

[0118]



wherein each symbol is as defined above.


[0119] In this method, the 2-aminobenzophenone derivative (IV) is reacted with the nitrile derivative (VI-1) to prepare the quinazoline derivative (I-2). The reaction of the compound (IV) with the compound (VI-1) is carried out in an appropriate solvent. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), ethers (e.g., dioxane, tetrahydrofuran, dimethoxyethane, etc.), N,N-dimethylformamide, dimethyl sulfoxide, chloroform, dichloromethane, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane, acetic acid and the like.

[0120] The reaction of the compound (IV) with the compound (VI-1) is carried out in the presence of an appropriate acid such as Lewis acids (e.g., aluminium chloride, zinc chloride, etc.), sulfuric acid, trifluoroacetic acid, p-toluenesulfonic acid and the like. The amount of the acid to be used is about 1 to 5 mol, preferably 1 to 2 mol per mol of the compound (IV). The reaction temperature is normally 20°C to 200°C, preferably about 30°C to 150°C. The reaction time is 0.5 to 20 hours, preferably 1 to 10 hours. The reaction may be carried out using an excess amount of the compound (VI-1) as the solvent.

[0121] The quinazoline compound (I-2) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method C

[0122]



wherein X^2 is an oxygen atom or sulfur atom, Z^2 is an optionally substituted hydrocarbon group or an optionally substituted heterocyclic group whose ring-constituting carbon atom is attached to X^2 .

[0123] Examples of the optionally substituted hydrocarbon group represented by Z^2 in the formulas (VII) and (I-4) include the same groups as those described above for R or Z^1 . Examples of the optionally substituted heterocyclic group whose ring-constituting carbon atom is attached to X^2 include the same groups as those described above for R or Z^1 .


[0124] In this method, the compound (I-3) is prepared according to the same manner as in the method A and then reacted with the compound (VIII) to prepare the compound (I-4). The reaction of the compound (IV) with the compound (V-2) can be carried out according to the same manner as in the method A. The reaction of the compound (I-3) with the compound (VII) is carried out in an appropriate solvent. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), ethers (e.g., dioxane, tetrahydrofuran, dimethoxyethane, etc.), alcohols (e.g., methanol, ethanol, propanol, etc.), ethyl acetate, acetonitrile, pyridine, N,N-dimethylformamide, dimethyl sulfoxide, chloroform, dichloromethane, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane, acetone, 2-butanone and the like. These solvents can be used alone or as a mixture thereof.

[0125] The reaction of the compound (I-3) with the compound (VII) is carried out in the presence of an appropriate base such as alkaline metal salts (e.g., sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, etc.), silver carbonate (Ag_2CO_3), amines (e.g., pyridine, triethylamine, N,N-dimethylaniline, etc.), sodium hydride, potassium hydride or the like. The amount of the base to be used is preferably about 1 to 5 mol per mol of the compound (I-3). The reaction temperature is normally -20°C to 150°C, preferably about -10°C to 100°C.

[0126] The quinoline derivatives (I-3) and (I-4) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method D

[0127]




wherein each symbol is as defined above.

[0128] In this method, the compound (I-5) is prepared according to the same manner as in the method B and then reacted with the compound (VII) to give the compound (I-6). The reaction of the compound (I-5) with the compound (VII) can be carried out according to the same manner as described for the reaction of the compound (I-3) with the compound (VII) in the method C.

[0129] The quinazoline derivatives (I-5) and (I-6) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method E

[0130]



wherein each symbol is as defined above.


[0131] In this method, the compound (I-7) is subjected to reduction to give the alcohol (I-8). This reduction can be carried out by per se known methods, for example, with a metal hydride, metal hydride complex, diborane or substituted borane, catalytic hydrogenation or the like. That is, this reaction is carried out by treating the compound (I-7) with a reducing agent. Examples of the reducing agent include metals and metal salts such as alkaline metal borohydride (e.g., sodium borohydride, lithium borohydride, etc.), metal hydride complexes (e.g., lithium aluminium hydride, etc.), metal hydrides (e.g., sodium hydride, etc.), organic tin compounds (e.g., triphenyltin hydride, etc.), nickel compounds, zinc compounds and the like; catalytic reducing agents using a transition metal catalyst (e.g., palladium, platinum, rhodium, etc.) and hydrogen; diborane and the like.

[0132] This reaction is carried out in an organic solvent which does not have a detrimental effect on the reaction. The solvent is appropriately selected depending upon the kind of the reducing agent. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), halogenated hydrocarbons (e.g., chloroform, carbon tetrachloride, dichloromethane, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane, etc.), ethers (e.g., diethyl ether, tetrahydrofuran, dioxane, etc.), alcohols (e.g., methanol, ethanol, propanol, isopropanol, 2-methoxyethanol, etc.), amides (e.g., N,N-dimethylformamide, etc.) and the like. These solvents can be used alone or as a mixture thereof. The reaction temperature is normally -20°C to 150°C , preferably about 0°C to 100°C . The reaction time is about 1 to 24 hours.

[0133] The quinoline derivative (I-8) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method F

[0134]



wherein q is an integer of 1 to 5, Y' is a nitrogen atom or $C-COOR^4$, and the other symbols are as defined above.

[0135] In this method, firstly, a compound of the formula (I-9) is reacted with an equivalent amount of triphenylphosphine to give a phosphonium salt of the formula (VIII). This reaction is carried out in a solvent. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), ethers (e.g., tetrahydrofuran, dioxane, dimethoxyethane, etc.), acetonitrile and the like. These solvents may be used alone or as a mixture thereof. The reaction temperature is 10°C to 200°C, preferably 30°C to 150°C. The reaction time is 0.5 to 50 hours.

[0136] Then the phosphonium salt (VIII) and aldehyde derivative (IX) are subjected to condensation reaction to give the compound (I-10). The condensation reaction of the compound (VIII) and the compound (IX) is carried out in an appropriate solvent in the presence of a base. Examples of the solvent include alcohols (e.g., methanol, ethanol, propanol, etc.), ethers (e.g., ethyl ether, dioxane, tetrahydrofuran, dimethoxyethane, etc.), aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), dichloromethane, 1,2-dichloroethane, *N,N*-dimethylformamide, dimethyl sulfoxide and the like. These solvents can be used alone or as a mixture thereof. Examples of the base include alkaline metal hydrides (e.g., sodium hydride, potassium hydride, etc.), alkoxides (e.g., sodium ethoxide, sodium methoxide, potassium ethoxide, potassium *tert*-butoxide, etc.), organic lithium compounds (e.g., methyl lithium, butyl lithium, phenyl lithium, etc.), sodium amide and the like. The amount of the base to be used is preferably about 1 to 1.5 mol per mol of the compound (VIII). The reaction temperature is normally -50°C to 100°C, preferably -20°C to 50°C. The reaction time is 0.5 to 20 hours.


[0137] The compound (I-10) can be obtained as a mixture of (E)- and (Z)-isomers with respect to the newly formed

double bond. By using each of the isomers after isolation or as a mixture of its isomers without isolation, the compound (I-10) is subjected to reduction to give the compound (I-11). This reduction is carried out according to conventional methods in a solvent in the presence of a catalyst such as palladium catalysts (e.g., palladium carbon, palladium black, etc.), platinum catalysts (e.g., platinum dioxide, etc.), Raney nickel or the like under an atmosphere of hydrogen. Examples of the solvent include alcohols (e.g., methanol, ethanol, propanol, etc.), ethers (e.g., ethyl ether, dioxane, tetrahydrofuran, dimethoxyethane, etc.), aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), dichloromethane, 1,2-dichloroethane, ethyl acetate, acetonitrile, acetone, 2-butanone, N,N-dimethylformamide, dimethyl sulfoxide and the like. These solvents may be used alone or as a mixture thereof. The hydrogen pressure is 1 to 150 atm, preferably 1 to 20 atm.

[0138] The quinoline or quinazoline derivatives (I-10) and (I-11) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method G

[0139]



wherein each symbol is as defined above.

[0140] In this method, firstly, the quinoline ester derivative (I-7) is subjected to hydrolysis reaction to give the carboxylic acid derivative (I-12). This hydrolysis reaction can be carried out according to conventional methods in the presence of an acid or base in a solvent. As the solvent, there can be used, for example, a mixture of water and an alcohol (e.g., methanol, ethanol, etc.), ether (e.g., tetrahydrofuran, dioxane, etc.), N,N-dimethylformamide, dimethyl sulfoxide, acetone or the like. Examples of the base include potassium carbonate, sodium carbonate, sodium methoxide, sodium ethoxide, potassium tert-butoxide, sodium hydroxide, potassium hydroxide, lithium hydroxide and the like. Examples of the acid include hydrochloric acid, sulfuric acid, acetic acid, hydrobromic acid and the like. The acid or base is preferably used in an excess amount (base: 1.2 to 10 equivalents, acid: 2 to 50 equivalents) based on the compound (I-7). The temperature of the reaction is normally -20°C to 150°C , preferably about -10°C to 100°C .


[0141] Then, the compound (I-12) is subjected to amidation to give the compound (I-13). This amidation is carried out by the reaction of the compound (I-12) with the compound (X). The condensation reaction of the compound (I-12) with the compound (X) is carried out by conventional techniques for peptide synthesis. The techniques for peptide synthesis may be by any known methods such as methods described in M. Bodansky and M. A. Ondetti, *Peptide Synthesis*, Interscience, New York (1966); F. M. Finn and K. Hofmann, *The Proteins*, Vol. 2, edited by H. Nenrath and R. L. Hill, Academic Press Inc., New York (1976); Nobuo Izumiya et al., *Basics and Experiments of Peptide Synthesis*, Maruzen K. K.

(1985), for example, azide method, chloride method, acid anhydride method, mixed anhydride method, DCC method, activated ester method, method using the Woodward reagent K, carbonyldiimidazole method, oxidation and reduction method, DCC/HONB method and method using DEPC (diethyl phosphorocyanidate). This condensation reaction can be carried out in a solvent. Examples of the solvent include anhydrous or hydrous dimethylformamide, dimethyl sulfoxide, pyridine, chloroform, dichloromethane, tetrahydrofuran, acetonitrile and the like. These solvents can be used alone or as a mixture thereof. The reaction temperature is normally about -20°C to about 50°C, preferably -10°C to 30°C. The reaction time is 1 to 100 hours, preferably 2 to 40 hours.

[0142] The quinoline derivatives (I-12) and (I-13) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method H

[0143]



wherein each symbol is as defined above.

[0144] The reaction of the compound (IV) with malonic acid ester derivative (XI) is carried out in the presence of a base according to per se known method. Examples of the base include organic bases such as trialkylamine (e.g., trimethylamine, triethylamine, etc.), picoline, N-methylpyrrolidine, N-methylmorpholine, 1,5-diazabicyclo[4.3.0]non-5-en, 1,4-diazabicyclo[2.2.2]non-5-en, 1,8-diazabicyclo[5.4.0]-7-undecene and the like. The amount of the base to be used is preferably 1 to 5 mol per mol of the compound (IV). The temperature of this reaction is normally 0°C to 200°C, preferably about 20°C to 150°C. Then the compound (XII) is chlorinated to give the compound (XIII). This method is car-


ried out by known methods, for example, by heating the compound (XII) with phosphorus pentachloride, phosphorus oxychloride, thionyl chloride or oxalyl chloride in a solvent or in the absence of a solvent.

[0145] The compound (XIII) is reacted with the compound (VII) to give the quinoline derivative (I-14). The reaction of the compound (XIII) with the compound (VII) is carried out in an appropriate solvent. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), ethers (e.g., dioxane, tetrahydrofuran, dimethoxyethane, etc.), alcohols (e.g., methanol, ethanol, propanol, etc.), ethyl acetate, acetonitrile, pyridine, N,N-dimethylformamide, dimethyl sulfoxide, chloroform, dichloromethane, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane, acetone, 2-butanone and the like. These solvents can be used alone or as a mixture thereof. The reaction of the compound (XIII) with the compound (VII) is carried out in the presence of a base such as alkaline metal salts (e.g., sodium hydroxide, potassium hydroxide, sodium carbonate, sodium bicarbonate, etc.), amines (e.g., pyridine, triethylamine, N,N-dimethylaniline, etc.), sodium hydride, potassium hydride or the like. The amount of the base to be used is preferably about 1 to 5 mol per mol of the compound (XIII). The reaction temperature is normally - 20°C to 150°C, preferably about -10°C to 100°C.

[0146] The quinoline derivative (I-14) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method I

[0147]




wherein r is 1 or 2, and the other symbols are as defined above.

[0148] In this method, the compound (I-15) obtained in Methods C or D is subjected to oxidation to give the compound (I-16). This oxidation is carried out according to conventional methods using an oxidizing agent such as m-chloroperoxybenzoic acid, hydrogen peroxide, peresters, sodium metaperiodate or the like. This oxidation is advantageously carried out in an organic solvent inert in the reaction conditions such as halogenated hydrocarbons (e.g., dichloromethane, chloroform, 1,2-dichloroethane, etc.), hydrocarbons (e.g., benzene, toluene, etc.), alcohols (e.g., methanol, ethanol, propanol, etc.) or the like. When the oxidizing agent is used in an equivalent amount or less based on the compound (I-15), the compound of the formula (I-16) wherein r is 1 is preferentially formed. When the oxidizing agent is used in excess of an equivalent amount, the compound of the formula (I-16) wherein r is 1 is further oxidized to give the compound of the formula (I-16) wherein r is 2. The reaction is carried out at room temperature or lower, preferably about -50°C to 20°C for 0.5 to 10 hours.

[0149] The quinoline derivative (I-16) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method J

[0150]




20 wherein each symbol is as defined above.

[0151] In this method, the compound (I-17) wherein X^2 is a sulfur atom that is obtained in Method H is subjected to oxidation to give the compound (I-18). This oxidation can be carried out according the same manner as in the method I.

[0152] The quinoline derivative (I-18) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method K

[0153]



wherein Q' is a leaving group and the other symbols are as defined above.

[0154] Examples of the leaving group represented by Q' include halogen atoms, preferably chlorine, bromine and iodine; hydroxyl groups activated by esterification such as organic sulfonic acid residues (e.g., p-toluenesulfonyloxy, methanesulfonyloxy, etc.), organic phosphoric acid residues (e.g., diphenylphosphoryloxy, dibenzylphosphoryloxy, dimethylphosphoryloxy, etc.) and the like.

[0155] In this method, firstly, the compound (I-19) is subjected to oxidation to give the compound (I-20). This oxidation is carried out according to the same manner as in the method I using an oxidizing agent such as m-chloroperbenzoic acid, hydrogen peroxide, peresters, sodium metaperiodate or the like. This oxidation is advantageously carried out in an organic solvent inert in the reaction conditions such as halogenated hydrocarbons (e.g., dichloromethane, chloroform, 1,2-dichloroethane, etc.), hydrocarbons (e.g., benzene, toluene, etc.), alcohols (e.g., methanol, ethanol, propanol, etc.) or the like. The amount of the oxidizing agent to be used is 1 to 5 mol, preferably 1 to 2 mol per mol of the compound (I-19). The reaction is carried out at 0°C to 120°C, preferably about -10°C to 100°C for normally 0.5 to 10 hours.


[0156] Then, the compound (I-20) is reacted with the compound (VII) according to the same manner as that described for the conversion from the compound (I-5) to the compound (I-6) in Method D.

[0157] The quinoline or quinazoline derivatives (I-20) and (I-21) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

[0158] The compound (I-21) wherein X² is a sulfur atom that is prepared by Method K can be converted to the corresponding sulfinyl or sulfonyl compound according to the same manner as that described in Method I.

Method L

[0159]




wherein each symbol is as defined above.

[0160] In this method, the compound (IV) is reacted with a tetronic acid derivative (XIV) to give the compound (II-1). This reaction is carried out according to the same manner as in the method A.

[0161] The quinoline derivative (II-1) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method M

[0162]



wherein each symbol is as defined above.


[0163] In this method, the compound (I-22) is reacted with the amine derivative (XV) to give the compound (III-1). The reaction of the compound (I-22) with the compound (XV) is carried out in an appropriate solvent. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), ethers (e.g., dioxane, tetrahydrofuran, dimethoxyethane, etc.), alcohols (e.g., methanol, ethanol, propanol, etc.), ethyl acetate, acetonitrile, pyridine, N,N-dimethylformamide, dimethyl sulfoxide, chloroform, dichloromethane, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane, acetone, 2-butanone and the like. These solvents can be used alone or as a mixture thereof. The reaction of the compound (I-22) with the compound (XV) is carried out in the presence of an appropriate base such as alkaline metal salts (e.g., sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, etc.), amines (e.g., pyridine, triethylamine, N,N-dimethylaniline, etc.), sodium hydride, potassium hydride or the like. The amount of the base to be used is preferably about 1 to 5 mol per mol of the compound (I-22). The reaction temperature

is normally -20°C to 150°C, preferably about -10°C to 100°C.

[0164] The quinoline derivative (III-1) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method N

[0165]



wherein each symbol is as defined above.

[0166] The reaction of the compound (I-23) with the compound (X) is carried out in an appropriate solvent. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), ethers (e.g., dioxane, tetrahydrofuran, dimethoxyethane, etc.), alcohols (e.g., methanol, ethanol, propanol, etc.), ethyl acetate, acetonitrile, pyridine, N,N-dimethylformamide, dimethyl sulfoxide, chloroform, dichloromethane, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane, acetone, 2-butanone and the like. These solvents can be used alone or as a mixture thereof. This reaction is carried out in the presence of an appropriate base such as alkaline metal salts (e.g., sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, etc.), amines (e.g., pyridine, triethylamine, N,N-dimethylaniline, etc.), sodium hydride, potassium hydride or the like. The amount of the base to be used is preferably about 1 to 5 mol per mol of the compound (I-23). The reaction temperature is normally -20°C to 150°C, preferably about -10°C to 100°C.


[0167] In this reaction, the compound (X) may be used as the base by using it in an excess amount.

[0168] The quinoline or quinazoline derivative (I-24) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

[0169] The quinoline derivative obtained by Methods L and M can be converted into the corresponding quinoline 1-oxide according to the oxidation method in Method K.

Method O

[0170]



wherein each symbol is as defined above.

[0171] In this method, the compound (XII) is converted into the mercapto derivative (XVI), followed by reaction with the compound (XVII) to give the compound (I-25). The reaction from the compound (XII) to the compound (XVI) can be carried out using a thiation reagent such as phosphorous pentasulfide (P_2S_5) or Lawesson's reagent or the like. Examples of the solvent include aromatic hydrocarbons (e.g., benzene, toluene, xylene, etc.), ethers (e.g., dioxane, tetrahydrofuran, dimethoxyethane, etc.), pyridine, chlorobenzene, dichlorobenzene, chloroform, dichloromethane, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane and the like. These solvents can be used alone or as a mixture thereof. The amount of the thiation reagent to be used is preferably about 1 to 5 mol per mol of the compound (XII). The reaction temperature is normally 0°C to 200°C , preferably about 10°C to 180°C . The reaction of the compound (XVI) with the compound (XVII) can be carried out according to the same manner as described for the reaction between the compounds (I-3) and (VII) in the method C.

[0172] The quinoline derivative (I-25) thus obtained can be isolated and purified by conventional separation and purification techniques such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, redistribution, chromatography and the like.

Method P

[0173] In this method, the compound (I) containing an isopropoxy group as the substituent of the ring A or ring B is treated with titanium tetrachloride, titanium trichloride, boron trichloride, tetrachlorosilane or the like to convert the isopropoxy group to a hydroxyl group, affording the corresponding compound containing a phenolic hydroxyl group as the substituent of the ring A or ring B.

[0174] This reaction is carried out in an appropriate solvent. Examples of the solvent include carbon tetrachloride, dichloromethane, chloroform, 1,2-dichloroethane, 1,1,2,2-tetrachloroethane, acetonitrile and the like. These solvents can be used alone or as mixtures thereof. The amount of the titanium tetrachloride, boron trichloride, tetrachlorosilane, etc., is 1 to 10 mol, preferably 1 to 6 mol per mol of the isopropoxy group. The reaction temperature is -50°C to 100°C , preferably -20°C to 80°C .

[0175] The following experiment illustrates the bone resorption inhibitory activity on the pharmaceutical composition of the present invention.

Experiment 1

Bone resorption inhibiting activity:

- 5 **[0176]** The bone resorption inhibiting activity was determined by the method of Raisz (J. Clin. Invest. 44, 103-116 (1965)). That is, ^{45}Ca (radioisotope of calcium in CaCl_2 solution)(50 μCi) was subcutaneously injected into a Sprague-Dawley rat of 18th day of pregnancy. On the next day, the abdomen was opened and a fetal rat was taken out sterilely. The left and right humeri (radii and ulnae) were removed from the body under a dissection microscope, and connective tissues and cartilages were removed as much as possible. Thus, bone culture samples were prepared. The bone was
- 10 incubated in a medium (0.6 ml) of BCJ_b Medium (Fitton-Jackson modification: GIBCO Laboratories, U.S.A.) containing 2 mg/ml of bovine serum albumin at 37°C for 24 hours in an atmosphere of 5% CO_2 in air. The bones were cultured for an additional 2 days in the above medium containing a final concentration of 1 $\mu\text{g/ml}$, 10 $\mu\text{g/ml}$ or 10 μM of the compound. This bone was cultivated for 2 days in the resulting medium. The radioactivity of ^{45}Ca in the medium and the radioactivity of ^{45}Ca in the bone were determined. The ratio (%) of ^{45}Ca released from the bone into the medium was
- 15 calculated according to the following equation.

$$\text{The ratio of } ^{45}\text{Ca} \text{ released from the bone into the medium (\%)} = \frac{^{45}\text{Ca counts in the medium}}{^{45}\text{Ca counts in the medium} + ^{45}\text{Ca counts in the bone}} \times 100$$

- 25 **[0177]** The bones from the same litter were cultured for 2 days by the same manner without addition of the compound, and used as the control. The mean \pm standard deviation of the values for five bones of each group was calculated. The ratio (%) of this value to the control value was calculated. The results are shown in Table 1 (In the tables hereinafter, Ex. No. indicates Example No.).

Table 1

Compound (Ex. No.)	Concentration	Bone resorption inhibitory activity (^{45}Ca release) (%) based on the control value)
28	10 $\mu\text{g/ml}$	49
28	1 $\mu\text{g/ml}$	51
29	10 $\mu\text{g/ml}$	47
41	10 $\mu\text{g/ml}$	45
41	1 $\mu\text{g/ml}$	67
53	10 $\mu\text{g/ml}$	59
57	10 $\mu\text{g/ml}$	59
68	10 $\mu\text{g/ml}$	86
78	10 $\mu\text{g/ml}$	66
81	10 $\mu\text{g/ml}$	55
87	10 $\mu\text{g/ml}$	59
89	10 $\mu\text{g/ml}$	50
89	1 $\mu\text{g/ml}$	78
131	10 $\mu\text{g/ml}$	79
134	10 $\mu\text{g/ml}$	73
137	1 $\mu\text{g/ml}$	54
139	10 $\mu\text{g/ml}$	61

Table 1 (continued)

Compound (Ex. No.)	Concentration	Bone resorption inhibitory activity (^{45}Ca release) (%) based on the control value)
142	10^{-5} M	66
148	10^{-5} M	56
159	10 $\mu\text{g/ml}$	39
166	10^{-5} M	34
181	10^{-5} M	45
189	10 $\mu\text{g/ml}$	58
211	10^{-5} M	28
237	10^{-5} M	40
243	10^{-5} M	50
263	10^{-5} M	45
264	10^{-5} M	51

[0178] As described above, the present invention provides a pharmaceutical composition comprising a quinoline or quinazoline derivative which has a direct effect on bones, exhibits excellent inhibitory activity of bone resorption and is useful as an agent for preventing or treating osteoporosis.

[0179] The following Examples and Reference examples further illustrate the present invention in detail but are not to be construed to limit the scope thereof.

Example 1

[0180] Conc. sulfuric acid (0.3 ml) was added to a mixture of 2-amino-3',4'-dimethoxy-4,5-ethylenedioxybenzophenone (6.5 g), ethyl 4-chloroacetoacetate (3.7 g) and acetic acid (60 ml), and the mixture was stirred at 100°C for 3 hours. The reaction mixture was concentrated under reduced pressure, and the residue was poured into water, made alkaline with 2N sodium hydroxide and extracted with chloroform. The chloroform layer was washed with water and dried over MgSO_4 . The solvent was evaporated under reduced pressure, and the residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (=7/3, v/v) gave ethyl 2-chloromethyl-4-(3,4-dimethoxyphenyl)-6,7-ethylenedioxyquinoline-3-carboxylate (5.5 g, 60%) which was then recrystallized from acetone.


Colorless prisms.
mp. 197-198°C.












Elemental Analysis:			
Calcd. for $\text{C}_{23}\text{H}_{22}\text{NO}_6\text{Cl}$	C,62.24;	H,5.00;	N,3.16
Found	C,61.95;	H,5.15;	N,3.01.

Examples 2 to 26


[0181] According to the same manner as that described in Example 1, the compounds in Tables 2 to 4 were obtained.













Table 2



Ex. No.	A ¹ , A ²		Yield (%)	mp (°C)	Recrystallization Solvent
2	6-Cl, H		61	105-106	Ethanol-Water
3	6-Cl, H		27	112-114	Methanol-Water
4	6-Cl, H		42	140-141	Ethyl acetate-Hexane
5	6-Cl, H		44	135-136	Ethyl acetate-Ethyl ether
6	6-CH₃, H		42	78-79	Ethyl acetate-Hexane
7	7-CH₃, H		40	125-126	Acetone-Ethyl ether
8	6-Br, H		58	108-109	Acetone-Isopropyl ether
9	6-CF₃, H		80	Note 1) Oil	_____
10	6, 7-(CH₃)₂		70	170-171	Ethyl acetate
11	6, 7-(CH₃)₂		42	119-120	Ethyl acetate-Hexane


Note 1) NMR (δ ppm) in CDCl₃: 0.92 (3H, t, J=7.2Hz), 4.06 (2H, q, J=7.2Hz), 5.03 (2H, s), 7.33-7.37 (2H, m), 7.50-7.55 (3H, m), 7.90-7.98 (2H, m), 8.26 (1H, d, J=9.4Hz).






Table 3

Ex. No.	A ¹ , A ²		Yield (%)	mp (°C)	Recrystallization Solvent
12	6, 7-(OCH ₂ CH ₂ O)		44	155-156	Acetone-Ethyl ether
13	6, 7-(CH ₃ O) ₂		23	153-155	Acetone-Ethyl ether
14	6, 7-(CH ₃ O) ₂		48	108-109	Ethyl ether
15	6, 7-(CH ₃ O) ₂		81	75-76	Isopropyl ether
16	6, 7-(CH ₃ O) ₂		53	146-147	Ethyl acetate-Hexane
17	6, 7-(CH ₃ O) ₂		50	151-153	Ethyl acetate-Hexane
18	6, 7-(CH ₃ O) ₂		53	160-161	Ethyl acetate-Hexane
19	6, 7-(CH ₃ O) ₂		35	126-127	Acetone-Ethyl ether
20	6, 7-(CH ₃ O) ₂		44	181-182 ¹⁾	Acetone-Ethyl ether
21	6, 7-(CH ₃ O) ₂		53	147-148	Acetone-Ethyl ether
22	6, 7-(CH ₃ O) ₂		44	134-135	Ethyl acetate-Hexane

Note 1) Methyl 2-chloromethyl-4-(3,4-dimethoxyphenyl)-
6,7-dimethoxyquinoline-3-carboxylate.

Table 4



Ex. No.	A ¹ , A ²		Yield (%)	mp (°C)	Recrystallization Solvent
23	6, 7-(CH ₃ O) ₂		64	211-212	Chloroform-Acetone
24	6, 7-(C ₂ H ₅ O) ₂		68	124-125	Ethyl acetate-Hexane
25	H, H		50	82-83	Ethyl acetate-Hexane
26	6-CH ₃		80	125-126	Ethanol

Example 27

[0182] Conc. sulfuric acid was added to a mixture of 2-amino-4,5,3',4'-tetramethoxybenzophenone, ethyl acetoacetate and acetic acid. The mixture was treated according to the same manner as that described in Example 1 to give ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-methylquinoline-3-carboxylate (83%) which was then recrystallized from ethanol.

Colorless prisms.
mp. 147-148°C.

Examples 28 to 49

[0183] According to the same manner as that described in Example 27, the compounds in Tables 5 to 9 were obtained.





Table 5

Ex. No.	A ¹ , A ²		R	R ⁴	Yield (%)	mp (°C)	Recrystallization Solvent
28	6, 7-(CH ₃ O) ₂		C ₂ H ₅	CH ₃	90	150-151	Ethanol
29	6, 7-(CH ₃ O) ₂		CH(CH ₃) ₂	CH ₃	40	160-161	Ethanol
30	6, 7-(CH ₃ O) ₂		CH ₃	C(CH ₃) ₃	32	139-141	I sopropyl ether - Hexane
31	6, 7-(CH ₃ O) ₂		C ₂ H ₅	CH ₃	76	110-111	Ethyl ether - I sopropyl ether
32	6, 7-(CH ₃ O) ₂		CH ₃	CH ₃	87	136-137	Ethanol

Table 6











Ex. No.	A ¹ , A ²		R	R ⁴	Yield (%)	mp (°C)	Recrystallization Solvent
33	6, 7-(CH ₃ O) ₂		CH ₃	CH ₃	77	129-130	E thanol
34	6, 7-(CH ₃ O) ₂		CH ₃	CH ₃	78	109-111	E thyl ether
35	6, 7-(OCH ₂ O)		CH ₃	CH ₃	45	155-156	E thanol
36	6, 7-(CH ₃ O) ₂		C ₂ H ₅	CH ₃	52	120-121	I sopropyl ether
37	6, 7-(CH ₃ O) ₂		CH ₃	CH ₃	75	154-155	I sopropyl ether

Table 7




Ex. No.	A ¹ , A ²		R	R ⁴	Yield (%)	mp (°C)	Recrystallization Solvent
38	6, 7-(CH ₃ O) ₂		CH ₃	C ₂ H ₅	63	197-199	E thanol
39	6-CH ₃ O		C ₂ H ₅	CH ₃	57	82-84	E thyl ether - Hexane
40	6-Cl		C ₂ H ₅	CH ₃	57	118-120	E thyl ether - Hexane
41	6-Cl		CH ₃	CH ₃	70	126-128	E thyl ether
42	6, 7-(CH ₃ O) ₂		C ₂ H ₅	CH ₃	45	102-103	E thyl ether - Hexane

Table 8



Ex. No.	A ¹ , A ²		R	R ⁴	Yield (%)	mp (°C)	Recrystallization Solvent
43	6-CH ₃		C ₂ H ₅	CH ₃	63	114-115	Ethyl ether - Hexane
44	6, 7-(CH ₃ O) ₂		C ₂ H ₅	CH ₃	73	136-137	Ethyl ether - Hexane
45	6-Cl		CH ₃	CH ₃	88	121-122	Ethyl ether - Hexane
46	6, 7-(CH ₃) ₂		C ₂ H ₅	CH ₃	68	127-128	I sopropyl ether
47	6-CH ₃		C ₂ H ₅	CH ₃	59	81-82	Ethyl ether

Table 9



Ex. No.	A ¹ , A ²		R	R ⁴	Yield (%)	mp (°C)	Recrystallization Solvent
48	6, 7-(CH ₃ O) ₂		CH ₃	C ₃ H ₇	79	153-155	Ethyl acetate - Isopropyl ether
49	6, 7-(CH ₃ O) ₂		CH ₃	C ₄ H ₉	53	119-120	Ethyl acetate - Hexane


Example 50

[0184] One drop of conc. sulfuric acid was added to a mixture of 2-amino-4,5,3',5'-tetramethoxy-4'-hydroxybenzophenone (0.333 g), tetronic acid (tetrahydrofuran-2,4-dione)(0.11 g) and acetic acid (10 ml), and the mixture was stirred at 100°C for 1.5 hours. The reaction mixture was concentrated under reduced pressure. The residual oil was poured into water, neutralized with an aqueous saturated sodium bicarbonate solution and extracted with chloroform. The chloroform layer was washed with water and dried over magnesium sulfate. Evaporation of the solvent gave 6,7-dimethoxy-9-(4-hydroxy-3,5-dimethoxyphenyl)furo[3,4-b]quinoline -1(3H)-one (0.349 g, 88%) which was then recrystallized from methanol. Pale yellow needles. mp. 247-248°C.

Examples 51 to 55

[0185] According to the same manner as that described in Example 50, the compounds in Table 10 were obtained.

Table 10



Ex. No.	A ¹	A ²		Yield (%)	mp (°C)	Recrystallization Solvent
51	CH ₃ O	CH ₃ O		90	241-242	Methanol - Chloroform
52	OCCH ₃ O			84	252-253	E thanol
53	CH ₃ O	CH ₃ O		38	210-211	E thanol
54	H	C1		89	210-212	Acetone
55	H	C1		95	254-255	Methanol

Example 56

[0186] A mixture of ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (3.0 g), m-chloroperbenzoic acid (85%, 2.3 g) and methanol (40 ml) was stirred for 2 hours under reflux. The solvent in the reaction mixture was evaporated under reduced pressure, and the residue was poured into chloroform. The chloroform layer was washed with water and dried over magnesium sulfate. The solvent was evaporated under reduced pressure. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (=6/4,

v/v) gave ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate 1-oxide (2.0 g, 65%) which was then recrystallized from acetone-isopropyl ether.

Colorless prisms.
mp. 193-194°C.

Elemental Analysis:

Calcd. for $C_{23}H_{24}NO_7Cl$	C,59.81;	H,5.24;	N,3.03
Found	C,59.69;	H,5.32;	N,3.05.

Example 57

[0187] Powdered aluminium chloride (6.7 g) was added to a mixture of 2-amino-4,5,3',4'-tetramethoxybenzophenone (8.0 g) and chloroacetonitrile (25 ml), and the mixture was stirred at 100°C for 2 hours. To the reaction mixture was poured into water, and the resulting mixture was extracted with chloroform. The chloroform layer was washed with water and dried over magnesium sulfate. The solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (=10/1, v/v) gave 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinazoline (4.9 g, 52%) which was then recrystallized from acetone.

Colorless prisms. mp. 183-184°C.

Example 58

[0188] A mixture of sodium iodide (1.68 g) and methyl ethyl ketone (15 ml) was stirred at 80°C for 1 hour, and then ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (2.0 g) was added thereto, and the resulting mixture was stirred at the same temperature for 12 hours. The insoluble solid was filtered off, and the filtrate was concentrated under reduced pressure. The residue was poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (1/1,v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-iodomethylquinoline-3-carboxylate (1.4 g, 58%) which was then recrystallized from ethyl acetate-hexane.

Colorless prisms.
mp. 170-171°C.

Elemental Analysis:

Calcd. for $C_{23}H_{24}NO_6I$	C,51.41;	H,4.50;	N,2.61
Found	C,51.25;	H,4.53;	N,2.58.

Example 59

[0189] A mixture of ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (3.0 g), 1-ethyl-2-mercaptoimidazole (1.0 g), potassium carbonate (1.1 g) and N,N-dimethylformamide (30 ml) was stirred at room temperature for 3 hours. The reaction mixture was poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (=3/2, v/v) gave ethyl 2-[(1-ethylimidazol-2-yl)thiomethyl]-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (2.8 g, 78%) which was then recrystallized from ethyl acetate-hexane.

Colorless prisms.

mp. 157-158°C.

5

Elemental Analysis:			
Calcd. for $C_{23}H_{31}N_3O_6S$	C, 62.55;	H, 5.81;	N, 7.82
Found	C, 62.55;	H, 5.84;	N, 7.79.

10

Example 60

[0190] m-Chloroperbenzoic acid (85%, 830 mg) was added in small portions under ice-cooling to a solution of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate (3.0 g) in dichloromethane (75 ml). The reaction mixture was stirred at room temperature for 2.5 hours, washed successively with 5% aqueous $NaHSO_3$ solution, saturated aqueous sodium bicarbonate solution and water, and dried over magnesium sulfate. The solvent was evaporated under reduced pressure. The residue was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/methanol (10/1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)sulfinylmethyl]quinoline-3-carboxylate (1.8 g, 58%) which was then recrystallized from acetone-ethyl ether.

20

Colorless prisms.
mp. 193-194°C.

25

Elemental Analysis:			
Calcd. for $C_{27}H_{29}N_3O_7S$	C, 60.10;	H, 5.42;	N, 7.79
Found	C, 59.80;	H, 5.60;	N, 7.51.

30

Example 61

[0191] According to the same manner as that described in Example 60, ethyl 2-[(2-benzimidazolyl)sulfinylmethyl]-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate was obtained and then recrystallized from acetone.

35

Colorless prisms.
mp. 160-161°C.

40

Elemental Analysis:			
Calcd. for $C_{30}H_{29}N_3O_7S$	C, 62.60;	H, 5.08;	N, 7.30
Found	C, 62.21;	H, 5.10;	N, 7.09.

45

Example 62

50

[0192] m-Chloroperbenzoic acid (85%, 2.5 g) was added in small portions under ice-cooling to a solution of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate (2.5 g) in dichloromethane (60 ml). The reaction mixture was stirred at room temperature for 4 hours, washed successively with 5% aqueous $NaHSO_3$ solution, saturated aqueous sodium bicarbonate solution and water, and dried over magnesium sulfate. The solvent was evaporated under reduced pressure. The residue was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/methanol (10/1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)sulfonylmethyl]quinoline-3-carboxylate (1.5 g, 58%) which was then recrystallized from acetone-ethyl ether.

55

Colorless prisms.
mp. 183-184°C.

Elemental Analysis:

Calcd. for $C_{27}H_{29}N_3O_8S$	C,58.37;	H,5.26;	N,7.56
Found	C,58.46;	H,5.24;	N,7.20.

Example 63

[0193] According to the same manner as that described in Example 62, ethyl 2-[(2-benzimidazolyl)sulfonylmethyl]-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate was obtained and then recrystallized from acetone-isopropyl ether.

Colorless prisms.
mp. 181-182°C.

Elemental Analysis:

Calcd. for $C_{30}H_{29}N_3O_8S$	C,60.90;	H,4.94;	N,7.10
Found	C,60.76;	H,4.86;	N,7.09.

Example 64

[0194] A solution of hydrogen chloride in ethanol (27%, 1.3 g) was added dropwise to a solution of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate (4.9 g) in ethanol (100 ml) at room temperature. About two thirds of the solvent was evaporated under reduced pressure, ethyl ether was added to the residue, and the resulting crystals were collected by filtration. The crystals were recrystallized from isopropanol to give ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate hydrochloride monohydrate (3.0 g, 55%).

Colorless prisms.
mp. 133-134°C.

Elemental Analysis:

Calcd. for $C_{27}H_{29}N_3O_6S \cdot HCl \cdot H_2O$	C,56.10;	H,5.58;	N,7.27
Found	C,55.84;	H,5.72;	N,7.16.

Example 65

[0195] According to the same manner as that described in Example 59, ethyl 6,7-dimethoxy-4-(2-methoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate (89%) was obtained as an oil.

[0196] NMR (δ ppm) in $CDCl_3$: 0.90 (3H,t, \underline{J} =7Hz), 3.34 (3H,s), 3.70 (3H,s), 3.74 (3H,s), 3.98 (2H,q, \underline{J} =7Hz), 4.03 (3H,s), 4.64 (2H,s), 6.66 (1H,s), 6.86 (1H,s), 7.01-7.16 (4H,m), 7.34 (1H,s), 7.45 (1H,double t, \underline{J} =8 and 2 Hz).

[0197] This oil was dissolved in ethanol (15 ml), and a solution of hydrogen chloride in ethanol (23%, 1.2 g) was added. Evaporation of the solvent under reduced pressure gave ethyl 6,7-dimethoxy-4-(2-methoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate hydrochloride (2.0 g) which was then recrystallized from ethanol-ethyl ether.

Pale yellow prisms.
mp. 180-181°C.

Elemental Analysis:			
Calcd. for $C_{26}H_{27}N_3O_5S \cdot HCl \cdot 1/2H_2O$	C, 57.93;	H, 5.42;	N, 7.80
Found	C, 58.05;	H, 5.32;	N, 7.72.

Example 66

[0198] According to the same manner as that described in Example 59, ethyl 6,7-dimethyl-4-(3,4-dimethylphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate (97%) was obtained as an oil.

NMR (δ ppm) in $CDCl_3$: 0.93 (3H, t, $J=7$ Hz), 2.31 (3H, s), 2.32 (3H, s), 2.35 (3H, s), 2.44 (3H, s), 3.42 (3H, s), 4.03 (2H, q, $J=7$ Hz), 4.61 (2H, s), 6.88 (1H, d, $J=1$ Hz), 7.03-7.10 (3H, m), 7.23 (1H, d, $J=8$ Hz), 7.35 (1H, s), 7.78 (1H, s).

[0199] This oil was dissolved in ethanol (10 ml), and a solution of hydrogen chloride in ethanol (23%, 0.584 g) was added. Evaporation of the solvent under reduced pressure gave ethyl 6,7-dimethyl-4-(3,4-dimethylphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate hydrochloride (1.1 g) which was then recrystallized from ethanol-ethyl ether.

Pale yellow prisms.
mp. 133-134°C.

Elemental Analysis:			
Calcd. for $C_{27}H_{29}N_3O_2S \cdot HCl \cdot 3/2H_2O$	C, 62.00;	H, 6.36;	N, 8.03
Found	C, 62.31;	H, 6.01;	N, 7.98.

Example 67


[0200] According to the same manner as that described in Example 59, ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate 1-oxide (69%) was obtained and then recrystallized from ethyl acetate-hexane.

Colorless prisms.
mp. 171-172°C.


Elemental Analysis:			
Calcd. for $C_{27}H_{29}N_3O_7S$	C, 60.10;	H, 5.42;	N, 7.79
Found	C, 60.29;	H, 5.53;	N, 7.49.

Examples 68 to 130

[0201] According to the same manner as that described in Example 59, the compounds in Tables 11 to 20 were obtained.

Table 11

Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
68	6-Cl, H			64	117-118	Ethyl acetate-Hexane
69	6-Cl, H			48	137-138	Acetone-Ethyl ether
70	6-Cl, H			71	120-121	Acetone-Isopropyl ether
71	6-Cl, H			55	190-191	Acetone-Isopropyl ether
72	6-CH ₃ , H			58	132-133	Ethyl acetate-Hexane

Table 12

















Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
73	7-CH ₃ , H			58	98-99	Ethyl acetate-Hexane
74	6-Br, H			69	129-130	Ethyl acetate-Hexane
75	6-CF ₃ , H			54	108-109	Ethyl acetate-Hexane
76	6, 7-(CH ₃) ₂			64	114-115	Ethyl acetate-Hexane
77	6, 7-(OCH ₂ CH ₂ O)			69	180-181	Acetone-Ethyl ether
78	6, 7-(OCH ₂ CH ₂ O)			60	120-121	Acetone
79	6, 7-(CH ₃ O) ₂			60	101-102	Acetone-Ethyl ether

Table 13

















Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
80	6, 7-(CH ₃ O) ₂			85	105-107	Ethyl ether
81	6, 7-(CH ₃ O) ₂			72	123-124	Ethyl ether
82	6, 7-(CH ₃ O) ₂			57	99-100	Ethyl acetate-Hexane
83	6, 7-(CH ₃ O) ₂			48	102-103	Isopropyl ether
84	6, 7-(CH ₃ O) ₂			80	117-118	Ethyl acetate-Hexane
85	6, 7-(CH ₃ O) ₂			74	132-133	Ethyl acetate-Hexane
86	6, 7-(CH ₃ O) ₂			46	134-135	Acetone-Ethyl ether

Table 14

















Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
87	6, 7- (CH ₃ O) ₂			81	145-146	Ethyl acetate-Hexane
88	6, 7- (CH ₃ O) ₂			77	147-148	Acetone-Isopropyl ether
89	6, 7- (CH ₃ O) ₂			84	149-150	Acetone-Ethyl ether
90	6, 7- (CH ₃ O) ₂			76	176-177	Acetone-Ethyl ether
91	6, 7- (CH ₃ O) ₂			65	111-112	Acetone-Ethyl ether
92	6, 7- (CH ₃ O) ₂			88	162-163	Acetone
93	6, 7- (CH ₃ O) ₂			77	185-186	Acetone

Table 15




















Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
94	6, 7-(CH ₃ O) ₂			90	165-166	Ethyl ether
95	6, 7-(CH ₃ O) ₂			83	152-153	Ethyl ether
96	6, 7-(CH ₃ O) ₂			86	174-176	Ethyl ether
97	6, 7-(CH ₃ O) ₂			80	184-185	Acetone
98	6, 7-(CH ₃ O) ₂			72	186-187	Ethyl acetate-Hexane
99	6, 7-(CH ₃ O) ₂			83	219-220	Ethyl acetate-Hexane
100	6, 7-(CH ₃ O) ₂			63	190-191	Ethyl acetate-Hexane

Table 16

Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
101	6, 7-(CH ₃ O) ₂			77	132-133	Ethyl acetate-Hexane
102	6, 7-(CH ₃ O) ₂			67	122-123	Ethyl acetate-Hexane
103	6, 7-(CH ₃ O) ₂			48	159-160	Ethyl acetate-Hexane
104	6, 7-(CH ₃ O) ₂			51	142-143	Ethyl acetate-Hexane
105	6, 7-(CH ₃ O) ₂			72	151-152	Ethyl acetate-Ethyl ether
106	6, 7-(CH ₃ O) ₂			64	188-189	Acetone-Isopropyl ether
107	6, 7-(CH ₃ O) ₂			82	188-190	Acetone-Isopropyl ether

Table 17




















Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
108	6, 7-(CH ₃ O) ₂			85	155-156	Methanol
109	6, 7-(CH ₃ O) ₂			77	173-174	Ethyl ether-Isopropyl ether
110	6, 7-(CH ₃ O) ₂			92	212-213	Ethyl ether-Isopropyl ether
111	6, 7-(CH ₃ O) ₂			72	118-120	Ethyl ether-Hexane
112	6, 7-(CH ₃ O) ₂			71	182-183	Acetone-Isopropyl ether
113	6, 7-(CH ₃ O) ₂			88	160-161	Ethyl acetate-Hexane
114	6, 7-(CH ₃ O) ₂			80	169-170	Ethyl ether

Table 18

Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
115	6, 7-(CH ₃ O) ₂			42	151-152	Acetone-Ethyl ether
116	6, 7-(CH ₃ O) ₂			36	167-168	Acetone-Ethyl ether
117	6, 7-(CH ₃ O) ₂			81	183-184	Ethyl acetate
118	6, 7-(CH ₃ O) ₂			71	235-237	Dichloromethane-Ethyl ether
119	6, 7-(CH ₃ O) ₂			89	198-199	Methanol
120	6, 7-(CH ₃ O) ₂			83	170-171	Acetone
121	6, 7-(CH ₃ O) ₂			88	110-112	Methanol

Table 19











Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
122	6, 7- (CH ₃ O) ₂			70	176-177	Ethyl acetate- Hexane
123	6, 7- (CH ₃ O) ₂			85	152-153	Acetone- Isopropyl ether
124	6, 7- (CH ₃ O) ₂			86	131-132	Acetone- Isopropyl ether
125	6, 7- (C ₂ H ₅ O) ₂			73	132-133	Ethyl acetate- Hexane

Table 20

Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
126	6, 7-(CH ₃ O) ₂			79	145-146	Ethyl acetate -Hexane
127	6, 7-(CH ₃ O) ₂			50	199-200	Dichloromethane -Ethyl ether
128	6, 7-(CH ₃ O) ₂			76	151-152	Ethyl acetate -Hexane
129	H, H			95	141-142	Ethyl acetate -Hexane
130	6-Cl, H			65	128-129	Ethyl acetate -Hexane

Example 131

[0202] A mixture of 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinazoline (4.5 g), 2-mercaptoethanol (1.13 g), potassium carbonate (2.8 g) and N,N-dimethylformamide (50 ml) was stirred at room temperature for 2 hours. The reaction mixture was poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate. Evaporation of the solvent gave 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(2-hydroxyethylthiomethyl)quinazoline (4.1 g, 82%) which was then recrystallized from ethanol. Colorless prisms. mp. 154-

155°C.

Examples 132 to 138

5 **[0203]** According to the same manner as that described in Example 131, the compounds in Table 21 were obtained.

10

15

20

25

30


35

40

45

50

55


Table 21

Ex. No.	A ¹ , A ²		Z ²	Yield (%)	mp (°C)	Recrystallization Solvent
132	6, 7-(CH ₃ O) ₂			77	143-144	Acetone
133	6, 7-(CH ₃ O) ₂		-CH ₂ COOCH ₃	83	138-139	Acetone
134	6, 7-(CH ₃ O) ₂			82	143-144	Acetone
135	6, 7-(CH ₃ O) ₂			68	143-144	Acetone
136	6, 7-(CH ₃ O) ₂			81	184-185	Acetone
137	6, 7-(CH ₃ O) ₂			80	195-196	Acetone -Isopropyl ether
138	6, 7-(CH ₃ O) ₂		-CH ₂	75	132-133	Ethyl ether

Examples 139 to 141

[0204] According to the same manner as that described in Example 60, the compounds in Table 22 were obtained.

Table 22



Ex. No.		Y	R	Yield (%)	mp (°C)	Recrystallization Solvent
139		N		83	126-127	Acetone -Isopropyl ether
140		C-COOC ₂ H ₅		58	152-153	Ethyl acetate -Hexane
141		C-COOC ₂ H ₅		59	168-169	Ethyl acetate -Hexane

Example 142

[0205] [6,7-Dimethoxy-4-(3,4-dimethoxyphenyl)-3-ethoxycarbonylquinolin-2-yl]methyltriphenylphosphonium chloride (17.4 g) was added at room temperature to a solution of sodium ethoxide in ethanol (prepared from Na (0.62 g) and ethanol (150 ml)). Then a solution of 2-formyl-1-methylimidazole (3.7 g) in ethanol (20 ml) was added dropwise. The mixture was stirred at room temperature for 3 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/methanol (100/1, v/v) gave ethyl (E)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)vinyl]quinoline-3-carboxylate (8.3 g, 67%) which was then recrystallized from ethyl acetate.

Colorless prisms. mp. 206-208°C.

[0206] The fractions eluted thereafter gave ethyl (Z)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)vinyl]quinoline-3-carboxylate (2.6 g, 21%) as an oil.

NMR (δ ppm) in CDCl₃: 0.96 (3H,t, \underline{J} =7Hz), 3.35 (3H,s), 3.78 (3H,s), 3.87 (3H,s), 3.96 (3H,s), 3.97 (3H,s), 3.98

(2H,q, \underline{J} =7Hz), 6.69 (1H,d, \underline{J} =12Hz), 6.8-7.1 (7H,m), 7.13 (1H,s).

[0207] Each of the ethyl (E)- and (Z)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)vinyl]quinoline-3-carboxylate was subjected to catalytic hydrogenation under an atmosphere of hydrogen at 1 atm in ethanol/tetrahydrofuran (1/1, v/v) in the presence of palladium-carbon (5%) to give ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)ethyl]quinoline-3-carboxylate which was then recrystallized from ethanol. Colorless prisms. mp. 147-148°C.

Example 143

[0208] [6,7-Dimethoxy-4-(3,4-dimethoxyphenyl)quinazolin-2-yl]methyltriphenylphosphonium chloride (9.1 g) was added at room temperature to a solution of sodium ethoxide in ethanol (prepared from Na (0.394 g) and ethanol (100 ml)). Then a solution of 2-formyl-1-methylimidazole (1.7 g) in ethanol (10 ml) was added dropwise. The mixture was stirred at room temperature for 3 hours, poured into water and extracted with chloroform. The chloroform layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/methanol (20/1, v/v) gave (E)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)vinyl]quinazoline (5.1 g, 82%) which was then recrystallized from ethanol-chloroform.

Colorless prisms. mp. 254-255°C.

Elemental Analysis:			
Calcd. for $C_{24}H_{24}N_4O_4 \cdot 3/2H_2O$	C,62.73;	H,5.92;	N,12.19
Found	C,62.62;	H,5.85;	N,11.90

The fractions eluted thereafter gave (Z)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)vinyl]quinazoline (0.61 g, 10%) which was then recrystallized from ethanol-chloroform.


Colorless plate crystals. mp. 180-181°C.

Elemental Analysis:			
Calcd. for $C_{24}H_{24}N_4O_4 \cdot 1/2H_2O$	C,65.29;	H,5.71;	N,12.69
Found	C,65.28;	H,5.66;	N,12.42

[0210] Each of the (E)- and (Z)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)vinyl]quinazoline was subjected to catalytic hydrogenation under an atmosphere of hydrogen at 1 atm in chloroform/ethyl acetate (1/1, v/v) in the presence of palladium-carbon (5%) to give ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)ethyl]quinazoline which was then recrystallized from ethyl acetate. Colorless prisms. mp. 170-171°C.

Examples 144 to 149

[0211] According to the same manner as that described in Example 142, the compounds in Table 23 were obtained.

Table 23

Ex. No.	A ¹ , A ²		Z ¹	q	mp (°C)	Recrystallization Solvent
144	6, 7-(CH ₃ O) ₂			1	183-184	Ethyl acetate
145	6, 7-(CH ₃ O) ₂			1	155-156	Ethyl acetate -Hexane
146	6, 7-(C ₂ H ₅ O) ₂			1	134-135	Ethyl acetate -Hexane
147	6, 7-(CH ₃ O) ₂			1	Note 1) 112-113	Ethyl acetate -Hexane
148	6, 7-(CH ₃ O) ₂			1	140-141	Ethyl acetate -Hexane
149	6, 7-(CH ₃ O) ₂			1	132-133	Ethyl acetate

Note 1) 1/2 hydrate

Example 150

[0212] Ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)ethyl]quinoline-3-carboxylate (9.0 g) was suspended in ethanol (40 ml), and ethanolic hydrogen chloride (22%, 10 g) was added. After stirring the mixture at

room temperature for 5 minutes, ethyl ether (150 ml) was added, and the resulting crystals were collected by filtration and recrystallized from ethanol-ethyl ether to give ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)ethyl]quinoline-3-carboxylate dihydrochloride (9.1 g).

Pale yellow prisms. mp. 158-160°C.

Elemental Analysis:

Calcd. for $C_{28}H_{31}N_3O_6 \cdot 2HCl \cdot 1/3C_2H_5OH \cdot 1/2H_2O$	C,57.11;	H,6.02;	N,6.97
Found	C,57.03;	H,6.15;	N,7.00

Example 151

[0213] [6,7-Dimethoxy-4-(3,4-dimethoxyphenyl)-3-ethoxycarbonylquinolin-2-yl]methyltriphenylphosphonium chloride (3.0 g) was added at room temperature to a solution of sodium ethoxide in ethanol (prepared from Na (0.13 g) and ethanol (45 ml)). Then 3-(1-methylimidazol-2-yl)propionaldehyde (0.787 g) was added. The mixture was stirred at room temperature for 3 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/methanol (30/1, v/v) gave ethyl (E)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)ethyl]vinyl]quinoline-3-carboxylate (0.36 g, 15%) as an oil.

NMR (δ ppm) in $CDCl_3$: 1.03 (3H,t, \underline{J} =7Hz), 2.7-3.0 (4H,m), 3.60 (3H,s), 3.79 (3H,s), 3.87 (3H,s), 3.97 (3H,s), 4.05 (3H,s), 4.09 (2H,q, \underline{J} =7Hz), 6.7-7.2 (8H,m), 7.43 (1H,s).

[0214] The fractions eluted thereafter gave ethyl (Z)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)ethyl]vinyl]quinoline-3-carboxylate (0.2 g, 8%) as an oil.

NMR (δ ppm) in $CDCl_3$: 1.02 (3H,t, \underline{J} =7Hz), 2.8-3.2 (4H,m), 3.58 (3H,s), 3.80 (3H,s), 3.88 (3H,s), 3.96 (3H,s), 4.05 (3H,s), 4.07 (2H,q, \underline{J} =7Hz), 6.08 (1H,dt, \underline{J} =7.4&11.4Hz), 6.6-7.0 (7H,m), 7.42 (1H,s).

[0215] A mixture of the ethyl (E)- and (Z)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[2-(1-methylimidazol-2-yl)ethyl]vinyl]quinoline-3-carboxylates was subjected to catalytic hydrogenation under an atmosphere of hydrogen at 1 atm in ethanol/tetrahydrofuran (1/4, v/v) in the presence of palladium-carbon (5%) and treated with ethanolic hydrogen chloride to give ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[4-(1-methylimidazol-2-yl)butyl]quinoline-3-carboxylate which was then recrystallized from chloroform-ethyl acetate.

Pale yellow crystals. mp. 180-183°C.

Elemental Analysis:

Calcd. for $C_{30}H_{35}N_3O_6 \cdot 2HCl \cdot H_2O$	C,57.69;	H,6.29;	N,6.73
Found	C,57.48;	H,6.09;	N,6.60

Example 152

[0216] A mixture of ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (1.5 g), 2-hydroxy-6-methylpyridine (0.4 g), potassium carbonate (0.511 g) and N,N-dimethylformamide (20 ml) was stirred at 120°C for 2 hours. The reaction mixture was poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate gave 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(2-methyl-6-pyridyl)oxymethyl]quinoline-3-carboxylate (0.79 g, 46%) which was then recrystallized from chloroform-hexane.

Yellow prisms. mp. 173-174°C.

Elemental Analysis:			
Calcd. for $C_{29}H_{30}N_2O_7$	C,67.17;	H,5.83;	N,5.40
Found	C,66.97;	H,6.02;	N,5.16

Example 153

[0217] A mixture of ethyl 2-iodomethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (9.0 g), 2-hydroxy-1-methylimidazole (1.8 g), silver (I) carbonate (Ag_2CO_3) (5.1 g) and benzene (100 ml) was stirred at 50°C for 18 hours. The insoluble solid was filtered off. The filtrate was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (5/1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methyl-2-imidazolyl)oxymethyl]quinoline-3-carboxylate (0.8 g, 9%) which was then recrystallized from ethyl acetate-hexane.

Colorless prisms. mp. 151-152°C.

Elemental Analysis:			
Calcd. for $C_{27}H_{29}N_3O_7$	C,63.90;	H,5.87;	N,8.28
Found	C,63.74;	H,5.87;	N,7.99

Example 154

[0218] A mixture of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate (0.6 g), 2N sodium hydroxide (1.7 ml) and ethanol (12 ml) was refluxed for 6 hours. The reaction mixture was concentrated in vacuo. The residue was diluted with water, washed with ethyl acetate, made acidic with 2N hydrochloric acid and extracted with ethyl acetate. The extract was washed with water, dried over magnesium sulfate and concentrated in vacuo to give crystals. The crystals were recrystallized from ethanol-ethyl ether to give 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylic acid (0.3 g, 53%) as colorless prisms. mp. 213-214°C.

Elemental Analysis:			
Calcd. for $C_{25}H_{25}N_3O_6S \cdot 1/2H_2O$	C,59.51;	H,5.19;	N,8.32
Found	C,59.38;	H,5.40;	N,7.93

Example 155

[0219] Oily sodium hydride (60%, 0.1 g) was added to a solution of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1H)-quinolone-3-carboxylate (0.5 g) in N,N-dimethylformamide (20 ml) under ice-cooling. After stirring at room temperature for 30 minutes, 2-bromopropane (0.5 ml) was added. The mixture was stirred at 60°C for 5 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate. Evaporation of the solvent gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-isopropoxyquinoline-3-carboxylate (0.388 g, 70%) which was then recrystallized from ethyl ether. mp. 142-143°C.

Example 156


[0220] Oily sodium hydride (60%, 0.033 g) was added to a solution of 2-hydroxymethyl-1-methylimidazole (0.085 g) in N,N-dimethylformamide (7 ml) under ice-cooling. After stirring at the same temperature for 10 minutes, ethyl 2-chloro-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (0.3 g) was added. The mixture was stirred under ice-cooling for 1 hour followed by stirring at room temperature for 20 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/methanol (30/1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1-methylimidzol-2-yl)methoxyquinoline-3-carboxylate (0.26 g, 73%) which was then recrystallized from ethyl acetate-hexane. mp. 209-210°C.

Example 157

[0221] Oily sodium hydride (60%, 0.05 g) was added to a mixture of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-mercaptoquinoline-3-carboxylate (0.2 g), 2-chloromethyl-1-methylimidazole (0.09 g) and N,N-dimethylformamide (8 ml) under ice-cooling. The mixture was stirred at the same temperature for 1 hour followed by stirring at room temperature for 18 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/methanol (30/1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[(1-methylimidzol-2-yl)methylthio]quinoline-3-carboxylate (0.214 g, 87%). mp. 141-142°C.

Example 158

[0222] A mixture of ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (0.58 g), 2-aminomethyl-1-methylimidazole hydrochloride (0.24 g), potassium carbonate (0.63 g) and N,N-dimethylformamide (15 ml) was stirred at room temperature for 2 hours. The mixture was poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform/methanol (50/1, v/v) gave 2,3-dihydro-6,7-dimethoxy-9-(3,4-dimethoxyphenyl)-2-(1-methylimidzol-2-ylmethyl)-1-oxo-1H-pyrro[3,4-b]quinoline (0.27 g, 43%) of the formula:



which was then recrystallized from ethyl acetate. mp. 245-246°C.

Example 159

[0223] Three drops of conc. sulfuric acid was added to a mixture of 2-amino-4,5,3',4'-tetramethoxybenzophenone (1.55 g), dimethyl acetonedicarboxylate (0.936 g) and acetic acid (30 ml), and the resulting mixture was stirred at 100°C for 2.5 hours. The reaction mixture was concentrated under reduced pressure, the residue was poured into water, neutralized with an aqueous saturated sodium bicarbonate solution and extracted with chloroform. The chloroform layer was washed with water dried over magnesium sulfate, and the solvent was evaporated under reduced pressure. The resulting crystals were recrystallized from acetone to give methyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-3-methoxycarbonylquinoline-2-acetate (1.41 g, 64%). Colorless prisms. mp. 170-171.

Example 160

[0224] According to the same manner as that described in Example 159, ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-

nyl)-3-ethoxycarbonylquinoline-2-acetate was obtained and then recrystallized from ethyl acetate-isopropyl ether.

Colorless prisms. mp. 146-147°C.

Example 161

[0225] Powdered aluminium chloride (0.21 g) was added to a mixture of 2-amino-5-chloro-3',4'-dimethoxybenzophenone (0.23 g) and methyl cyanoacetate (5 ml), and the resulting mixture was stirred at 100°C for 2.5 hours. The reaction mixture was poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with hexane/ethyl acetate (4/1, v/v) gave methyl 6-chloro-4-(3,4-dimethoxyphenyl)quinazoline-2-acetate (0.161 g, 55%) which was then recrystallized from isopropyl ether. Colorless needles. mp. 144-145°C.

Example 162

[0226] According to the same manner as that described in Example 161, methyl 6-chloro-4-phenylquinazoline-2-acetate was obtained and recrystallized from isopropyl ether. Colorless needles. mp. 122-123°C.

Example 163

[0227] According to the same manner as that described in Example 161, methyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinazoline-2-acetate was obtained and recrystallized from isopropyl ether. Colorless needles. mp. 152-153°C.

Example 164

[0228] A solution of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-3-ethoxycarbonylquinoline-2-acetate (5.8 g) in tetrahydrofuran (100 ml) was added dropwise to a suspension of lithium aluminium hydride (0.455 g) in tetrahydrofuran (50 ml) at 0°C. After stirring the reaction mixture at 0°C for 1 hour, water (2.5 ml) was added dropwise, and the mixture was stirred for additional 30 minutes. The insoluble solid was filtered off, and the filtrate was concentrated under reduced pressure. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (1/1, v/v) gave ethyl 2-(2-hydroxyethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (1.75 g, 33%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 150-151°C.

Example 165

[0229] Phosphorus tribromide (PBr₃) (1.0 g) was added dropwise to a solution of ethyl 2-(2-hydroxyethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (1.7 g) in benzene (50 ml) at room temperature. The mixture was stirred at 80°C for 1 hour, poured into ice-water, neutralized with an aqueous saturated sodium bicarbonate solution and extracted with chloroform. The chloroform layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (1/1, v/v) gave ethyl 2-(2-bromoethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (0.49 g, 26%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 132-133°C.


Example 166

[0230] Oily sodium hydride (60%, 0.323 g) was added to a solution of 2-ethylimidazole (0.776 g) in N,N-dimethylformamide (30 ml), and the mixture was stirred at room temperature for 15 minutes. Then ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (3.0 g) was added. The mixture was stirred at 80°C for 1 hour and poured into water, and the resulting crystals were collected by filtration and recrystallized from ethanol to give ethyl 2-(2-ethylimidazol-1-ylmethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (2.5 g, 74%). Colorless prisms. mp. 163-164°C.

Examples 167 to 176


[0231] According to the same manner as that described in Example 166, the compounds in Tables 24 and 25 were obtained.

Table 24



Ex. No.		Y		mp (°C)	Recrystallization Solvent
167		C-COOC ₂ H ₅		208-209	Dichloromethane—Hexane
168		C-COOC ₂ H ₅		177-178	Ethyl acetate—Hexane
169		C-COOC ₂ H ₅		134-135	Ethanol
170		C-COOC ₂ H ₅		200-201	Ethyl acetate—Hexane
171		C-COOC ₂ H ₅		148-149	Ethyl acetate—Hexane

Table 25



Ex. No.		Y		mp (°C)	Recrystallization Solvent
172		C-COOC ₂ H ₅		157-158	Ethanol
173		N		184-185	Dichloromethane-Ethyl ether
174		N		223-224	Dichloromethane-Ethyl ether
175		N		188-189	Ethyl acetate-Hexane
176		N		198-199	Dichloromethane-Ethyl ether

Example 177


[0232] Oily sodium hydride (60%, 0.044 g) was added to a solution of imidazole (0.075 g) in N,N-dimethylformamide (5 ml), and the mixture was stirred at room temperature for 15 minutes. Then ethyl 2-(2-bromoethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (0.46 g) was added. The mixture was stirred at 80°C for 1 hour, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/methanol (10/1, v/v) gave ethyl 2-[2-(1-imidazolyl)ethyl]-6,7-

dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (0.295 g, 66%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 173-174°C.

Examples 178 to 180

[0233] According to the same manner as that described in Example 177, the compounds in Tables 26 were obtained.

Table 26



Ex. No.	B	Y	-N P	mp (°C)	Recrystallization Solvent
178		C-COOC ₂ H ₅		209-210	Dichloromethane - Hexane
179		C-COOC ₂ H ₅		198-199	Ethanol
180		C-COOC ₂ H ₅		124-125	Ethyl acetate - Hexane

Example 181

[0234] Oily sodium hydride (60%, 0.323 g) was added to a solution of 1H-1,2,4-triazole (0.558 g) in N,N-dimethylformamide (30 ml), and the mixture was stirred at room temperature for 15 minutes. Then ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (3.0 g) was added. The mixture was stirred at 80°C for 1 hour, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was subjected to column chromatography on silica gel. The fractions firstly eluted with chloroform/methanol (40/1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (1.7 g, 53%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 176-177°C.

Example 182

[0235] The fractions eluted thereafter of the column chromatography in Example 181 gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-4-ylmethyl)quinoline-3-carboxylate (0.07 g, 2%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 226-227°C.

Example 183

[0236] According to the same manner as that described in Example 181, ethyl 6,7-dimethoxy-4-(4-methoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 150-151°C.

Example 184

[0237] The fractions eluted thereafter of the column chromatography in Example 183 gave ethyl 6,7-dimethoxy-4-(4-methoxyphenyl)-2-(1,2,4-triazol-4-ylmethyl)quinoline-3-carboxylate which was then recrystallized from ethyl acetate-hexane. Colorless needles. mp. 218-219°C.

Example 185

[0238] According to the same manner as that described in Example 181, 6,7-dimethoxy-4-(4-methoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinazoline which was then recrystallized from dichloromethane-ethyl ether. Colorless prisms. mp. 206-207°C.

Example 186

[0239] The fractions eluted thereafter of the column chromatography in Example 185 gave 6,7-dimethoxy-4-(4-methoxyphenyl)-2-(1,2,4-triazol-4-ylmethyl)quinazoline which was then recrystallized from dichloromethane-ethyl ether. Colorless needles. mp. 212-213°C.

Example 187

[0240] A mixture of methyl 2-ethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (0.5 g), 4N sodium hydroxide (20 ml) and methanol was stirred for 14 hours under reflux, poured into water, made acidic and extracted with chloroform. The chloroform layer was washed with water and dried over magnesium sulfate. Evaporation of the solvent gave 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-ethylquinoline-3-carboxylic acid (0.343 g, 71%) which was then recrystallized from acetone-methanol. mp. 264-267°C.

Example 188

[0241] One drop of N,N-dimethylformamide was added to a mixture of 2-ethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylic acid (0.2 g), oxalyl chloride (0.05 ml) and tetrahydrofuran (10 ml), and the resulting mixture was stirred at room temperature for 1 hour and concentrated under reduced pressure. The residue was dissolved in tetrahydrofuran (10 ml). The solution was added dropwise to a mixture of diethyl 2-aminoethylphosphonate (0.195 g), triethylamine (0.1 ml) and tetrahydrofuran (10 ml). The reaction mixture was stirred at room temperature for 1 hour and then under reflux for 3 hours, poured into water and extracted with chloroform. The chloroform layer was washed successively with 2N hydrochloric acid, an aqueous saturated sodium bicarbonate solution and water, and dried over mag-

nesium sulfate. Evaporation of the solvent gave N-(2-diethoxyphosphorylethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-ethylquinoline-3-carboxamide (0.21 g, 75%) which was then recrystallized from ethyl ether. mp. 157-158°C.

5 Example 189

[0242] A solution of methyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-methylquinoline-3-carboxylate (0.4 g) in tetrahydrofuran (10 ml) was added dropwise to a suspension of lithium aluminium hydride (LiAlH_4) (0.114 g) in tetrahydrofuran (10 ml) at room temperature. The mixture was stirred at room temperature for 1.5 hours, and water was added. The resulting mixture was extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, the solvent was evaporated, and the resulting crystals were collected by filtration and recrystallized from methanol to give 3-hydroxymethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-methylquinoline (0.326 g, 88%). mp. 200-201°C.

15 Example 190

[0243] According to the same manner as that described in Example 189, 2-ethyl-3-hydroxymethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline was obtained and recrystallized from acetone-isopropyl ether. mp. 192-193°C.

20 Example 191

[0244] m-Chloroperbenzoic acid (80%, 1.4 g) was added under ice-cooling to a solution of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1-methylimidazol-2-ylmethylthio)quinoline-3-carboxylate (3.0 g) in dichloromethane (90 ml), and the mixture was stirred at the same temperature for 3 hours. The reaction mixture was washed successively with an aqueous saturated NaHSO_3 solution, aqueous saturated sodium bicarbonate solution and water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/dichloromethane (4:1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1-methylimidazol-2-ylmethylsulfinyl)quinoline-3-carboxylate (2.17 g, 70%). mp. 151-152°C.

30 Example 192

[0245] The fractions eluted thereafter of the column chromatography in Example 191 gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1-methylimidazol-2-ylmethylsulfonyl)quinoline-3-carboxylate (0.47 g, 14%). mp. 132-133°C.

35 Example 193

[0246] A mixture of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-methylquinoline-3-carboxylate (411 mg), N-bromosuccinimide (214mg), 2,2-azobis(isobutyronitrile) (10 mg) and carbon tetrachloride (10 ml) was stirred under reflux for 5 hours. The reaction mixture was washed with water, dried over magnesium sulfate. The solvent was evaporated under reduced pressure, and the residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (10/1, v/v) gave ethyl 2-bromomethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (285 mg, 58%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 135-136°C.

45

Elemental Analysis:			
Calcd. for $\text{C}_{23}\text{H}_{24}\text{NO}_6\text{Br}$	C,56.34;	H,4.93;	N,2.86
Found	C,55.98;	H,5.23;	N,2.62

50

Example 194

[0247] According to the same manner as that described in Example 193, propyl 2-bromomethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (48%) was obtained and recrystallized from ethyl acetate-isopropyl ether. Colorless prisms. mp. 144-145°C

Elemental Analysis:			
Calcd. for C ₂₄ H ₂₆ NO ₆ Br	C,57.15;	H,5.20;	N,2.78
Found	C,56.75;	H,5.30;	N,2.68

Example 195

[0248] According to the same manner as that described in Example 193, butyl 2-bromomethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (56%) was obtained and recrystallized from ethyl acetate-ethyl ether. Colorless prisms. mp. 160-161°C.

Elemental Analysis:			
Calcd. for C ₂₅ H ₂₈ NO ₆ Br	C,57.92;	H,5.44;	N,2.70
Found	C,57.96;	H,5.53;	N,2.50

Example 196

[0249] According to the same manner as that described in Example 1, ethyl 2-chloromethyl-6,7-dimethoxy-4-(4-methoxy-3-propoxyphenyl)quinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 126-128°C

Example 197

[0250] Methanol (15 ml) was added dropwise under reflux to a mixture of methyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinazoline-2-acetate (4.0 g), sodium borohydride (1.9 g) and tetrahydrofuran (80 ml). The mixture was stirred under reflux for 2 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water, dried over magnesium sulfate. The solvent was evaporated under reduced pressure to give 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(2-hydroxyethyl)quinazoline (3.0 g, 81%) which was then recrystallized from ethyl acetate. Colorless needle crystals. mp. 165-166°C.


Example 198

[0251] According to the same manner as that described in Example 165, 2-(2-bromoethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinazoline (56%) was obtained and recrystallized from ethyl acetate. Colorless needles. mp. 166-167°C.

Examples 199 to 205

[0252] According to the same manner as that described in Example 1, the compounds in Table 27 were obtained.

Table 27




Ex. No.	A¹	A²	B¹	B²	Yield (%)	mp (°C)	Recrystallization Solvent
199	CH₃O	CH₃O	Cl	Cl	57	159-160	Ethyl acetate-hexane
200	(CH₃)₂CHO	CH₃O	CH₃O	CH₃O	66	138-140	Ethyl acetate-hexane
201	CH₃O	(CH₃)₂CHO	CH₃O	CH₃O	48	125-126	Ethyl acetate-hexane
202	CH₃O	CH₃O	(CH₃)₂CHO	CH₃O	50	126-127	Ethanol
203	CH₃O	CH₃O	CH₃O	(CH₃)₂CHO	48	149-150	Ethanol
204	CH₃O	CH₃O	(CH₃)₂CHO	(CH₃)₂CHO	48	118-119	Ethyl acetate-hexane
205	CH₃O	(CH₃)₂CHO	(CH₃)₂CHO	(CH₃)₂CHO	60	99-100	Ethyl acetate-hexane

Examples 206 to 214

[0253] According to the same manner as that described in Example 181, the compounds in Table 28 were obtained.

Table 28



Ex. No.	A¹	A²	B¹	B²	W	Yield (%)	mp (°C)	Recrystallization Solvent
206	CH ₃ O	CH ₃ O	H	Cl	CH	65	156-157	Ethyl acetate-Hexane
207	CH ₃ O	CH ₃ O	Cl	Cl	CH	64	183-184	Ethyl acetate-Hexane
208	CH ₃ O	CH ₃ O	Cl	Cl	N	48	160-161	Ethyl acetate-Hexane
209	(CH ₃) ₂ CHO	CH ₃ O	CH ₃ O	CH ₃ O	N	58	154-155	Ethyl acetate-Hexane
210	CH ₃ O	(CH ₃) ₂ CHO	CH ₃ O	CH ₃ O	N	62	-1)	
211	CH ₃ O	CH ₃ O	(CH ₃) ₂ CHO	CH ₃ O	N	65	183-185	Ethyl acetate-Hexane
212	CH ₃ O	CH ₃ O	CH ₃ O	(CH ₃) ₂ CHO	N	75	165-166	Ethyl acetate-Hexane
213	CH ₃ O	CH ₃ O	(CH ₃) ₂ CHO	(CH ₃) ₂ CHO	N	50	134-135	
214	CH ₃ O	(CH ₃) ₂ CHO	(CH ₃) ₂ CHO	(CH ₃) ₂ CHO	N	66	Oil ²⁾	

1) Amorphous solid. NMR (δ ppm in CDCl_3): 0.87
 (3H,t,J=7.2Hz), 1.33 (6H,d,J=6.0Hz), 3.85 (3H,s), 3.93
 (2H,q,J=7.2Hz), 3.96 (3H,s), 4.02 (3H,s), 4.43 (1H,m), 5.68
 (1H,d,J=14.8Hz), 5.77 (1H,d,J=14.8Hz), 6.82-7.01 (4H,m), 7.41
 (1H,s), 7.93 (1H,s), 8.27 (1H,s).

2) NMR (δ ppm in CDCl_3): 0.84 (3H,t,J=7.2Hz), 1.26-1.45
 (18H,m), 3.93 (2H,q,J=7.0Hz), 4.02 (3H,s), 4.21 (1H,m), 4.51
 (1H,m), 4.56 (1H,m), 5.73 (2H,s), 6.80-6.92 (3H,m), 7.01
 (1H,d,J=8.2Hz), 7.41 (1H,s), 7.93 (1H,s), 8.27 (1H,s).

Example 215


[0254] Titanium tetrachloride (TiCl_4) (125 mg) was added at 0°C to a solution of ethyl 6,7-dimethoxy-4-(4-isopropoxy-3-methoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (55.6 mg) in dichloromethane (2 ml), and the mixture was stirred at the same temperature for 6 hours. The reaction mixture was poured into water and extracted with chloroform. The chloroform layer was washed with a saturated aqueous sodium carbonate solution and water, and dried over magnesium sulfate. The solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/chloroform (3/2, v/v) gave ethyl 6,7-dimethoxy-4-(4-hydroxy-3-methoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (24.5 mg, 48%) which was then recrystallized from ethyl acetate-hexane. mp. 176-178°C.

NMR (δ ppm in CDCl_3): 0.88 (3H,t,J=7.2Hz), 3.80 (3H,s), 3.88 (3H,s), 3.96 (2H,q,J=7.2Hz), 4.05 (3H,s), 5.73 (2H,s), 5.80 (1H, broad s), 6.80-7.06 (4H,m), 7.42 (1H,s), 7.94 (1H,s), 8.27 (1H,s).

Examples 216 to 218

[0255] According to the same manner as that described in Example 215, the compounds in Table 29 were obtained.

Table 29



Ex. No.	A ¹	A ²	B ¹	B ²	Yield (%)	mp (°C)	Recrystallization Solvent
216	H ₂ O	CH ₃ O	CH ₃ O	CH ₃ O	35	165-166 ¹⁾	
217	CH ₃ O	H ₂ O	CH ₃ O	CH ₃ O	38	215-216 ²⁾	
218	CH ₃ O	CH ₃ O	H ₂ O	CH ₃ O	62	232-233	Dichloromethane-hexane

1) NMR (δ ppm in CDCl_3): 0.88 (3H,t,J=7.2Hz), 3.84
 (3H,s), 3.86 (3H,s), 3.95 (2H,q,J=7.2Hz), 3.97 (3H,s), 5.73
 (2H,s), 6.88–7.01 (5H,m), 7.52 (1H,s), 7.94 (1H,s), 8.37
 (1H,s).

2) NMR (δ ppm in CDCl_3): 0.86 (3H,t,J=7.0Hz), 3.85
 (3H,s), 3.94 (2H,q,J=7.0Hz), 3.98 (3H,s), 4.07 (3H,s), 5.73
 (2H,s), 6.20 (1H,broad), 6.82–6.98 (3H,m), 7.08 (1H,s), 7.42
 (1H,s), 7.93 (1H,s), 8.27 (1H,s).

Example 219

[0256] Titanium tetrachloride (TiCl_4)(288 mg) was added at 0°C to a solution of ethyl 4-(3,4-diisopropoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (116.0 mg) in dichloromethane (2.5 ml), and the mixture was stirred at the same temperature for 6 hours. The reaction mixture was poured into water and extracted with chloroform. The chloroform layer was washed with a saturated aqueous sodium bicarbonate solution and water, and dried over magnesium sulfate. The solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/chloroform (7/3, v/v) gave ethyl 4-(3,4-dihydroxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (20.0 mg, 21%). mp. 122–124°C.

NMR (δ ppm in CDCl_3): 0.78 (3H,t,J=7.0Hz), 3.78 (3H,s), 3.86 (2H,q,J=7.0Hz), 4.00 (3H,s), 5.71 (2H,s), 6.60 (1H, broad s), 6.68–6.79 (2H,m), 6.92 (1H,s), 6.97 (1H,d,J=8.0Hz), 7.37 (1H,s), 7.95 (1H,s), 8.35 (1H,s), 8.70 (1H, broad s).

Example 220

[0257] Titanium tetrachloride (TiCl_4)(316 mg) was added at 0°C to a solution of ethyl 4-(3,4-diisopropoxyphenyl)-6-isopropoxy-7-methoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (96.0 mg) in dichloromethane (1.0 ml), and the mixture was stirred at the same temperature for 10 hours. The reaction mixture was poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with a saturated aqueous sodium bicarbonate solution and water, and dried over magnesium sulfate. The solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/methanol (10/1, v/v) gave ethyl 4-(3,4-dihydroxyphenyl)-6-hydroxy-7-methoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (19.0 mg, 26%). mp. 264–266°C.

NMR (δ ppm in CDCl_3): 0.88 (3H,t,J=7.0Hz), 3.93 (2H,q,J=7.0Hz), 3.94 (3H,s), 5.63 (2H,s), 6.52 (1H,dd,J=8.2&2.2Hz), 6.67 (1H,d,J=2.2Hz), 6.85 (1H,d, J=8.2Hz), 6.98 (1H,s), 7.29 (1H,s), 7.94 (1H,s), 8.57 (1H,s), 9.17 (1H,s), 9.21 (1H,s), 10.00 (1H,s).

Example 221

[0258] Conc. sulfuric acid (0.03 ml) was added to a mixture of 2-amino-4,5,3',4'-tetramethoxybenzophenone (453 mg), ethyl 6-(1-imidazolyl)-3-oxohexanoate (320 mg) and acetic acid (5 ml), and the mixture was stirred at 100°C for 2 hours. The reaction mixture was concentrated under reduced pressure. The residue was poured into water, and the mixture was made alkaline with 2N sodium hydroxide and extracted with chloroform. The chloroform layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform/methanol (50/1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[3-(1-imidazolyl)propyl]quinoline-3-carboxylate (310.0 mg, 43%) which was then recrystallized

from ethyl acetate-hexane. Colorless prisms. mp. 164-165°C.

Example 222

- 5 **[0259]** According to the same manner as that described in Example 221, ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[3-(1,2,4-triazol-1-yl)propyl]quinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 141-142°C.

Example 223

- 10 **[0260]** According to the same manner as that described in Example 221, ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-[4-(1-imidazolyl)butyl]quinoline-3-carboxylate was obtained and recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 119-120°C.

15 Example 224

- [0261]** A mixture of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (3.0 g), 2N sodium hydroxide (15.6 ml) and ethanol (50 ml) was stirred under reflux for 8 hours. The reaction mixture was ice-cooled, adjusted to pH 5 with 2N hydrochloric acid and concentrated under reduced pressure. The residue was dissolved in ethanol, the insoluble materials were filtered off, and the filtrate was concentrated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform/methanol (4/1, v/v) gave 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylic acid (1.3 g, 46%) which was then recrystallized from dichloromethane-ethanol. Colorless prisms. mp. 270-271°C (decomposition).

25 Example 225

- [0262]** Oily sodium hydride (60%, 0.156 g) was added to a solution of 1H-1,2,4-triazole (0.27 g) in N,N-dimethylformamide (DMF) (20 ml), and the mixture was stirred at room temperature for 15 minutes. Then ethyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate 1-oxide (1.5 g) was added, and the mixture was stirred at 80°C for 45 minutes. The reaction mixture was poured into water and extracted with dichloromethane. The dichloromethane layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform/methanol (30/1, v/v) gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate 1-oxide (0.8 g, 50%) which was then recrystallized from dichloromethane-hexane. Colorless prisms. mp. 221-222°C.

35 Example 226

- [0263]** According to the same manner as that described in 181, ethyl 6,7-dimethoxy-4-(3-propoxy-4-methoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 127-128°C.

Example 227

- [0264]** In the column chromatography in Example 226, the fractions eluted thereafter gave ethyl 6,7-dimethoxy-4-(3-propoxy-4-methoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate which was then recrystallized from ethanol. Colorless needles. mp. 154-155°C.

Example 228

- 50 **[0265]** According to the same manner as that described in 181, ethyl 4-(3,4-dimethoxyphenyl)-6,7-ethylenedioxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless needles. mp. 138-140°C.

Example 229

- 55 **[0266]** In the column chromatography in Example 228, the fractions eluted thereafter gave ethyl 4-(3,4-dimethoxyphenyl)-6,7-ethylenedioxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate which was then recrystallized from ethanol. Colorless needles. mp. 237-239°C.

Example 230

[0267] According to the same manner as that described in 181, ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,3-triazol-1-ylmethyl)quinoline-3-carboxylate was obtained and recrystallized from ethanol-dichloromethane. Colorless prisms. mp. 195-196°C.

Elemental Analysis:			
Calcd. for $C_{25}H_{26}N_4O_6 \cdot 1/4C_2H_5OH$	C,62.50;	H,5.66;	N,11.43
Found	C,62.29;	H,5.53;	N,11.30

Example 231

[0268] In the column chromatography in Example 230, the fractions eluted thereafter gave ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,3-triazol-2-ylmethyl)quinoline-3-carboxylate which was then recrystallized from ethanol-dichloromethane. Colorless needles. mp. 163-164°C.

Elemental Analysis:			
Calcd. for $C_{25}H_{26}N_4O_6 \cdot 1/2C_2H_5OH$	C,62.27;	H,5.83;	N,11.17
Found	C,61.98;	H,5.69;	N,11.10

Example 232

[0269] In the column chromatography in Example 211, the fractions eluted thereafter gave ethyl 6,7-dimethoxy-4-(3-isopropoxy-4-methoxyphenyl)-2-(1,2,4-triazol-4-ylmethyl)quinoline-3-carboxylate which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 170-171°C.

Example 233

[0270] In the column chromatography in Example 212, the fractions eluted thereafter gave ethyl 6,7-dimethoxy-4-(4-isopropoxy-3-methoxyphenyl)-2-(1,2,4-triazol-4-ylmethyl)quinoline-3-carboxylate which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 178-179°C.

Example 234

[0271] Oily sodium hydride (60%, 0.117 g) was added to a solution of 2-hydroxypyridine (0.277 g) in N,N-dimethylformamide (DMF)(10 ml), and the mixture was stirred at room temperature for 15 minutes. To this solution was added ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-iodomethylquinoline-3-carboxylate (1.2 g). The mixture was stirred at room temperature for 8 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (10/1, v/v) gave ethyl 2-(1,2-dihydro-2-oxopyridine-1-ylmethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (0.64 g, 57%) which was then recrystallized from ethanol. Colorless prisms. mp. 154-156°C.

Example 235

[0272] According to the same manner as that described in Example 177, 2-[2-(1-imidazolyl)ethyl]-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinazoline was obtained and recrystallized from ethyl acetate. Colorless prisms. mp. 147-148°C.

Example 236

[0273] According to the same manner as that described in Example 166, ethyl 2-(benzimidazol-1-ylmethyl)-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate was obtained by the reaction of ethyl 2-bromomethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate with benzimidazole, and recrystallized from ethanol. Colorless prisms. mp. 99-100°C.

Example 237

[0274] According to the same manner as that described in Example 181, methyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate was obtained by the reaction of methyl 2-chloromethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate with 1H-1,2,4-triazole, and recrystallized from ethanol. Colorless prisms. mp. 218-220°C.

Example 238

[0275] According to the same manner as that described in Example 166, propyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(imidazol-1-ylmethyl)quinoline-3-carboxylate was obtained by the reaction of propyl 2-bromomethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate with imidazole, and recrystallized from ethanol. Colorless prisms. mp. 166-168°C.

Example 239


[0276] According to the same manner as that described in Example 166, butyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-(imidazol-1-ylmethyl)quinoline-3-carboxylate was obtained by the reaction of butyl 2-bromomethyl-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate with imidazole, and recrystallized from ethanol. Colorless prisms. mp. 140-141°C.

Example 240

[0277] According to the same manner as that described in Example 181, ethyl 6-chloro-4-phenyl-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate was obtained by the reaction of ethyl 6-chloro-2-chloromethyl-4-phenylquinoline-3-carboxylate with 1H-1,2,4-triazole, and recrystallized from ethanol. Colorless prisms. mp. 114-116°C.

Examples 241 to 248

[0278] According to the same manner as that described in Example 166, the compounds in Table 30 were obtained.

Table 30

Ex. No.	A ¹ , A ²			Yield (%)	mp (°C)	Recrystallization Solvent
2 4 1	6-Cl, H			34	112-114	Ethanol
2 4 2	6-CH ₃ , H			30	121-123	Ethanol
2 4 3	6, 7-(CH ₃) ₂			40	133-135	Ethanol
2 4 4	6, 7-(CH ₃ O) ₂			57	143-144	Ethanol
2 4 5	6, 7-(CH ₃ O) ₂			43	139-141	Ethyl acetate-Hexane
2 4 6	6, 7-(OCH ₂ CH ₂ O)			68	154-156	Ethanol
2 4 7	6, 7-(CH ₃ O) ₂			70	143-144	Ethanol-Hexane
2 4 8	6, 7-(CH ₃ O) ₂			75	160-161	Dichloromethane-Isopropyl ether

Example 249

[0279] A solution of hydrogen chloride in ethanol (23%, 0.172 g) was added dropwise at room temperature to a suspension of ethyl 6,7-dimethoxy-4-(3-isopropoxy-4-methoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (0.5 g) in ethanol (10 ml)-dichloromethane (2 ml). The mixture was stirred at room temperature for 15 minutes and concentrated under reduced pressure. The residue was treated with isopropyl ether, and the solid was collected by filtration

and recrystallized from ethanol to give ethyl 6,7-dimethoxy-4-(3-isopropoxy-4-methoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate hydrochloride (0.211 g, 39%). Yellow crystals. mp. 93-95°C.

Example 250

[0280] Oily sodium hydride (60%, 0.27 g) was added to a solution of morpholine (0.537 g) in N,N-dimethylformamide (20 ml), and the mixture was stirred at room temperature for 15 minutes. Then ethyl 2-chloromethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (2.5 g) was added. The resulting mixture was stirred at 100°C for 2 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The crystals were collected by filtration and then recrystallized from ethanol to give ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-morpholinomethylquinoline-3-carboxylate (1.9 g, 68%). Colorless prisms. mp. 146-147°C.

Example 251

[0281] A mixture of ethyl 2-chloromethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (1.0 g), piperazine (1.15 g) and methanol (15 ml) was stirred at room temperature for 36 hours. The reaction mixture was concentrated under reduced pressure, and 6N HCl (30 ml) was added to the residue. The resulting mixture was washed with dichloromethane. The aqueous layer was neutralized with 2N NaOH and extracted with dichloromethane. The dichloromethane layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The crystals were collected by filtration and recrystallized from dichloromethane-hexane to give ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-piperazinomethylquinoline-3-carboxylate (0.43 g, 39%). Colorless prisms. mp. 192-193°C.

Example 252

[0282] Oily sodium hydride (60%, 0.753 g) was added to a solution of thiomorpholine (1.8 g) in N,N-dimethylformamide (50 ml), and the mixture was stirred at room temperature for 15 minutes. Then ethyl 2-chloromethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (6.0 g) was added. The resulting mixture was stirred at room temperature overnight, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/ethyl acetate (7/3, v/v) gave ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-thiomorpholinomethylquinoline-3-carboxylate (1.4 g, 42%) which was then recrystallized from ethanol. Colorless prisms. mp. 148-149°C.

Example 253

[0283] Oily sodium hydride (60%, 0.466 g) was added to a solution of N-methylhomopiperazine (1.23 g) in N,N-dimethylformamide (40 ml), and the mixture was stirred at room temperature for 15 minutes. Then ethyl 2-chloromethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (4.0 g) was added. The resulting mixture was stirred at 100°C for 3.5 hours, poured into water and extracted with ethyl acetate. The ethyl acetate layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform/methanol (5/1, v/v) gave ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[(4-methylhomopiperazino)methyl]quinoline-3-carboxylate (1.0 g, 22%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 157-159°C.

Example 254

[0284] A mixture of ethyl 2-chloromethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (4.0 g), diethylamine (1.28 g) and methanol (45 ml) was stirred at room temperature for 4 days. The reaction mixture was poured into water and extracted with dichloromethane. The dichloromethane layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate gave ethyl 2-(N,N-diethylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (0.51 g, 18%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 130-131°C.

Example 255

[0285] A mixture of ethyl 2-chloromethyl-6,7-dimethoxy-4-(4-methoxyphenyl)quinoline-3-carboxylate (2.0 g), morpho-

line (2.5 g) and methanol (30 ml) was stirred at room temperature for 17 hours. The reaction mixture was concentrated under reduced pressure, and dichloromethane was added to the residue. The resulting mixture was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The crystals were collected by filtration and recrystallized from ethanol to give ethyl 6,7-dimethoxy-4-(4-methoxyphenyl)-2-morpholinomethylquinoline-3-carboxylate (1.65 g, 74%). Colorless prisms. mp. 165-166°C.

Example 256

[0286] According to the same manner as that described in Example 255, ethyl 6,7-dimethoxy-4-(4-methoxyphenyl)-2-piperidinomethylquinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 129-130°C.

Example 257

[0287] A mixture of ethyl 2-chloromethyl-6,7-dimethoxy-4-(2-methoxyphenyl)quinoline-3-carboxylate (1.0 g), morpholine (1.25 g) and ethanol (13 ml) was stirred at room temperature for 3 days. The crystals were collected by filtration and recrystallized from ethyl acetate-hexane to give ethyl 6,7-dimethoxy-4-(2-methoxyphenyl)-2-morpholinomethylquinoline-3-carboxylate (0.76 g, 68%). Colorless prisms. mp. 153-154°C.

Example 258

[0288] According to the same manner as that described in Example 255, ethyl 4-(3,4-dimethoxyphenyl)-6,7-ethylenedioxy-2-morpholinomethylquinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless needles. mp. 174-175°C.

Example 259

[0289] According to the same manner as that described in Example 255, ethyl 4-(3,4-dimethoxyphenyl)-6,7-ethylenedioxy-2-piperidinomethylquinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 158-160°C.

Example 260

[0290] According to the same manner as that described in Example 255, ethyl 4-(3,4-dimethoxyphenyl)-6,7-diethoxy-2-morpholinomethylquinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 147-148°C.

Example 261

[0291] According to the same manner as that described in Example 255, ethyl 4-(3,4-dimethoxyphenyl)-6,7-diethoxy-2-piperidinomethylquinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 154-155°C.

Example 262

[0292] According to the same manner as that described in Example 255, ethyl 6-chloro-2-morpholinomethyl-4-phenylquinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 161-163°C.

Example 263

[0293] A mixture of ethyl 2-chloromethyl-4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (3.0 g), morpholine (2.76 g) and ethanol (50 ml)-dichloromethane (5 ml) was stirred at room temperature overnight. The reaction mixture was concentrated under reduced pressure, and dichloromethane (50 ml) was added to the residue. The dichloromethane layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform-ethyl acetate (1/1, v/v) gave ethyl 4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxy-2-morpholinomethylquinoline-3-carboxylate as an oil. The oil was dissolved in ethanol (20 ml), a solution of hydrogen chloride in ethanol (23%, 1.05 g) was added, and the mixture was stirred at room temperature for 10 minutes. The solvent was evaporated under reduced pressure, and

the crystals were collected by filtration and recrystallised from ethanol-ether to give ethyl 4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxy-2-morpholinomethylquinoline-3-carboxylate hydrochloride (1.62 g, 45%). Colorless crystals. mp. 185-188°C.

Elemental Analysis:			
Calcd. for $C_{29}H_{37}N_2O_7Cl \cdot H_2O$	C,60.15;	H,6.79;	N,4.84
Found	C,60.51;	H,6.58;	N,4.73

Example 264

[0294] According to the same manner as that described in Example 263, ethyl 4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxy-2-piperidinomethylquinoline-3-carboxylate hydrochloride was obtained and recrystallized from ethanol-ether. Colorless crystals. mp. 207-210°C.

Elemental Analysis:			
Calcd. for $C_{30}H_{39}N_2O_6Cl \cdot 1/2H_2O$	C,63.43;	H,7.10;	N,4.93
Found	C,63.15;	H,7.02;	N,4.80

Example 265

[0295] According to the same manner as that described in Example 263, ethyl 2-(N,N-dipentylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate dihydrochloride was obtained and recrystallized from dichloromethane-ethyl acetate. Yellow powder. mp. 93-95°C.

Elemental Analysis:			
Calcd. for $C_{33}H_{48}N_2O_6Cl_2 \cdot 3/2H_2O$	C,59.45;	H,7.71;	N,4.20
Found	C,59.58;	H,7.88;	N,4.14

Example 266

[0296] According to the same manner as that described in Example 250, ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-piperidinomethylquinoline-3-carboxylate was obtained and recrystallized from ethanol. Colorless prisms. mp. 148-149°C.

Example 267

[0297] According to the same manner as that described in Example 250, ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-pyrrolidinomethylquinoline-3-carboxylate was obtained and recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 139-140°C.

Example 268

[0298] A mixture of propyl 2-bromomethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (1.5 g), piperidine (1.27 g) and dichloromethane (30 ml) was stirred at room temperature for 2 days. The reaction mixture was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform-ethyl acetate (10/1, v/v) gave propyl 4-(3,4-

dimethoxyphenyl)-6,7-dimethoxy-2-piperidinomethylquinoline-3-carboxylate (0.8 g, 53%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 129-131°C.

Example 269

5

[0299] According to the same manner as that described in Example 268, butyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-piperidinomethylquinoline-3-carboxylate was obtained and recrystallized from ethyl acetate-hexane. Colorless needles. mp. 154-155°C.

10 Examples 270 to 277

[0300] According the same manner as that described in Example 254, the compounds in Table 31 were obtained.

15

20

25

30

35


40

45

50

55

Table 3 1



Ex. No.	A ¹ , A ²	R	Yield (%)	mp (°C)	Recrystallization Solvent
2 7 0	6, 7-(CH ₃ O) ₂	CH ₃	65	137-138	Ethyl acetate-hexane
2 7 1	6, 7-(CH ₃ O) ₂	C ₃ H ₇	47	92-93	Ethanol-hexane
2 7 2	6, 7-(CH ₃ O) ₂	C ₄ H ₉	50	90-92	Ethanol-hexane
2 7 3	6, 7-(CH ₃ O) ₂	(CH ₃) ₂ CH	20	134-135	Ethanol-hexane
2 7 4	6, 7-(CH ₃ O) ₂	HOCH ₂ CH ₂	66	117-119 ¹⁾	Ethyl acetate-hexane
2 7 5	6, 7-(CH ₃ O) ₂	cyclohexyl	14	174-175 ²⁾	Ethyl acetate-hexane
2 7 6	6, 7-(CH ₃ O) ₂	C ₆ H ₅ CH ₂	36	56-58 ³⁾	
2 7 7	6, 7-(C ₂ H ₅ O) ₂	C ₂ H ₅	44	86-88	Ethyl acetate-hexane

1) 1/2 Hydrate.

2) 1/4 Hydrate.

3) Amorphous solid. NMR (δ ppm in CDCl_3): 0.97

(3H,t,J=7Hz), 3.59 (4H,s), 3.78 (3H,s), 3.88 (3H,s), 3.97
(3H,s), 3.99 (2H,q,J=7Hz), 4.04 (3H,s), 4.13 (2H,s), 6.84–7.02
(4H,m), 7.10–7.40 (10H,m), 7.42 (1H,s).

Example 278

[0301] A mixture of 2-chloromethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinazoline (2.0 g), piperidine (2.27 g) and ethanol (40 ml)-dichloromethane (10 ml) was stirred at room temperature for 3 days. The reaction mixture was concentrated under reduced pressure, and dichloromethane (50 ml) was added to the residue. The dichloromethane layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform-ethyl acetate (1/1, v/v) gave 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-piperidinomethylquinazoline (1.46 g, 65%) which was then recrystallized from ethyl acetate-hexane. Colorless needles. mp. 130-132°C.

Example 279

[0302] According to the same manner as that described in Example 278, 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-morpholinomethylquinazoline was obtained and recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 148-150°C.

Example 280

[0303] According to the same manner as that described in Example 278, 2-(N,N-diethylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinazoline was obtained and recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 111-113°C.

Example 281

[0304] According to the same manner as that described in Example 278, ethyl 2-(N-ethyl-N-propylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate was obtained by the reaction of ethyl 2-bromomethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate with N-ethyl-N-propylamine, and recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 105-106°C.

Example 282

[0305] A mixture of 2-(2-bromoethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinazoline (0.486 g), diethylamine (0.41 g) and dichloromethane (10 ml) was stirred under reflux for 6 hours. The reaction mixture was poured into water and extracted with dichloromethane. The dichloromethane layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform-ethyl acetate (1/1, v/v) gave 2-[2-(N,N-diethylamino)ethyl]-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinazoline (0.040 g, 8%) which was then recrystallized from ethyl acetate-hexane. Colorless prisms. mp. 164-166°C.


Example 283

[0306] A mixture of ethyl 2-chloromethyl-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (3.0 g), N-methyl-N-cyclohexylamine (2.28 g) and ethanol (45 ml) was stirred under reflux for 6 hours. The reaction mixture was poured into water and extracted with dichloromethane. The dichloromethane layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residue was subjected to column chromatography on silica gel. The fractions eluted with chloroform gave ethyl 2-(N-cyclohexyl-N-methylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate (2.80 g, 80%) which was then recrystallized from ethanol. Colorless prisms. mp. 172-174°C.

Examples 284 to 290

[0307] According to the same manner as that described in Example 283, the compounds in Table 32 were obtained.

Table 3 2



Ex. No.	R ¹	R ²	Yield (%)	mp (°C)	Recrystallization Solvent
2 8 4	C ₆ H ₅ CH ₂ CH ₂	C ₆ H ₅ CH ₂	65	135-137	Ethanol
2 8 5	C ₄ H ₉	CH ₃	64	143-144	Ethanol
2 8 6	(CH ₃) ₂ CHCH ₂	CH ₃	44	121-123	Ethyl acetate-Hexane
2 8 7	C ₄ H ₉	C ₂ H ₅	58	113-114	Ethyl acetate-Hexane
2 8 8	(C ₂ H ₅)(CH ₃)CH	C ₃ H ₇	39	129-131	Ethyl acetate-Hexane
2 8 9	(CH ₃) ₃ C	C ₂ H ₅	51	120-121	Ethyl acetate-Hexane
2 9 0	CH ₃	C ₂ H ₅	63	139-140	Ethyl acetate-Hexane

Example 291

[0308] According to the same manner as that described in Example 263, ethyl 2-(N,N-diethylaminomethyl)-4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate dihydrochloride was obtained by the reaction of ethyl

2-chloromethyl-4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate with diethylamine, and recrystallized from ethyl acetate-ether. Yellow powder. mp. 122-124°C.

Elemental Analysis:			
Calcd. for $C_{29}H_{40}N_2O_6Cl_2 \cdot 1/2H_2O$	C,58.78;	H,6.97;	N,4.73
Found	C,58.84;	H,7.00;	N,4.69

Example 292

[0309] According to the same manner as that described in Example 282, ethyl 6-chloro-4-(4-chlorophenyl)-2-(N,N-diethylaminomethyl)quinoline-3-carboxylate was obtained by the reaction of ethyl 6-chloro-2-chloromethyl-4-(4-chlorophenyl)-quinoline-3-carboxylate with diethylamine, and recrystallized from ethanol. Colorless prisms. mp. 132-133°C.

Example 293

[0310] According to the same manner as that described in Example 282, ethyl 6-chloro-2-(N,N-diethylaminomethyl)-4-phenylquinoline-3-carboxylate was obtained by the reaction of ethyl 6-chloro-2-chloromethyl-4-phenylquinoline-3-carboxylate with diethylamine, and recrystallized from ethanol. Colorless prisms. mp. 107-108°C.

Reference Example 1

[0311] A mixture of 2-amino-4,5,3',4'-tetramethoxybenzophenone (10.0 g), diethyl malonate (6.0 g) and 1,8-diazabicyclo[5.4.0]-7-undecene (DBU)(0.396 g) was stirred at 180°C for 10 minutes. After cooling, ethanol was added to the reaction mixture, and the crystals were collected by filtration to give ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2(1H)-quinolone-3-carboxylate (12.0 g, 91%) which was then recrystallized from chloroform-acetone. mp. 273-276°C.

Reference Example 2

[0312] A mixture of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2(1H)-quinolone-3-carboxylate (6.8 g), Lawesson's reagent (8.0 g) and toluene (250 ml) was stirred under reflux for 18 hours. After cooling, the precipitated crystals were collected by filtration to give ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2-mercaptoquinoline-3-carboxylate (5.0 g, 70%) which was then recrystallized from acetone. mp. 265-266°C.

Reference Example 3

[0313] A mixture of ethyl 6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-2(1H)-quinolone-3-carboxylate (5.0 g) and phosphorus oxychloride ($POCl_3$)(6.3 ml) was stirred at 100 to 110°C for 80 minutes. The reaction mixture was concentrated under reduced pressure. The residue was poured into water, neutralized with an aqueous saturated sodium bicarbonate solution and extracted with chloroform. The chloroform layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform gave ethyl 2-chloro-6,7-dimethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (3.8 g, 73%) which was then recrystallized from ethyl acetate-hexane. mp. 168-169°C.

Reference Example 4

[0314] A mixture of ethyl 2-chloromethyl-6,7-diethoxy-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate (7.1 g), triphenylphosphine (3.9 g) and toluene (70 ml) was stirred under reflux for 2 hours. After cooling, the resulting solid was collected by filtration to give [6,7-diethoxy-4-(3,4-dimethoxyphenyl)-3-ethoxycarbonylquinolin-2-yl]methyltriphenylphosphonium chloride (9.6 g, 87%). mp. 172-174°C (decomposition).

Reference Example 5

[0315] According to the same manner as that described in Reference Example 4, [6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-3-ethoxycarbonylquinolin-2-yl]methyltriphenylphosphonium chloride was obtained. mp. 200-202°C (decomposition).

sition).

Reference Example 6

5 **[0316]** According to the same manner as that described in Reference Example 4, [6,7-dimethoxy-4-(4-methoxyphenyl)-3-ethoxycarbonylquinolin-2-yl]methyltriphenylphosphonium chloride was obtained. mp. 178-180°C (decomposition).

Reference Example 7

10 **[0317]** According to the same manner as that described in Reference Example 4, [6,7-dimethoxy-4-(3,4-dimethoxyphenyl)-quinazolin-2-yl]methyltriphenylphosphonium chloride was obtained. mp. 208-210°C (decomposition).

Reference Example 8

15 **[0318]** A mixture of benzyl 4-bromobutyrate (23.7 g), imidazole (8.1 g), potassium carbonate (14.0 g) and acetone (400 ml) was stirred under reflux for 6 hours. After cooling the mixture to room temperature, the insoluble material was filtered off, and the filtrate was concentrated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with ethyl acetate/methanol (20/1, v/v) gave benzyl 4-(1-imidazolyl)butyrate (7.3 g, 33%) as an oil.

20 NMR (δ ppm in CDCl_3): 2.11 (2H,m), 2.34 (2H,t,J=6.8Hz), 3.99 (2H,t,J=6.8Hz), 5.12 (2H,s), 6.87 (1H,s), 7.05 (1H,s), 7.30-7.40 (5H,m).

Reference Example 9

25 **[0319]** According to the same manner as that described in Reference Example 8, benzyl 4-(1,2,4-triazol-1-yl)butyrate was obtained as an oil in 88% yield.

30 NMR (δ ppm in CDCl_3): 2.14-2.42 (4H,m), 4.24 (2H,t,J=6.4Hz), 5.13 (2H,s), 7.30-7.43 (5H,m), 7.94 (1H,s), 7.99 (1H,s).

Reference Example 10

35 **[0320]** According to the same manner as that described in Reference Example 8, benzyl 5-(1-imidazolyl)valerate was obtained as an oil by the reaction of benzyl 5-bromovalerate with imidazole.

NMR (δ ppm in CDCl_3): 1.55-1.90 (4H,m), 2.38 (2H,t,J=6.8Hz), 3.93 (2H,t,J=7.0Hz), 5.11 (2H,s), 6.87 (1H,s), 7.05 (1H,s), 7.25-7.50 (5H,m), 7.94 (1H,s), 7.99 (1H,s).

Reference Example 11

40 **[0321]** A mixture of benzyl 4-(1-imidazolyl)butyrate (7.4 g), palladium-carbon (5%) (1.0 g) and ethanol (400 ml) was subjected to catalytic hydrogenation at room temperature and 1 atm. The catalyst was filtered off, the filtrate was concentrated under reduced pressure, and the crystals were recrystallized from ethanol to give 4-(1-imidazolyl)butyric acid (3.4 g, 75%). Colorless prisms. mp. 125-126°C.

Reference Example 12

50 **[0322]** According to the same manner as that described in Reference Example 11, benzyl 4-(1,2,4-triazol-1-yl)butyrate was subjected to catalytic hydrogenation to give 4-(1,2,4-triazol-1-yl)butyric acid which was then recrystallized from ethanol. Colorless prisms. mp. 137-138°C.

Reference Example 13

55 **[0323]** According to the same manner as that described in Reference Example 11, benzyl 5-(1-imidazolyl)valerate was subjected to catalytic hydrogenation to give 5-(1-imidazolyl)valeric acid which was then recrystallized from ethanol. Colorless prisms. mp. 157-158°C.

Reference Example 14

[0324] To a suspension of 4-(1-imidazolyl)butyric acid (0.5 g) in tetrahydrofuran (35 ml) was added 1,1'-carbonyldiimidazole (0.578 g) was added, and the mixture was stirred at room temperature for 6 hours. Then malonic acid monoethyl ester magnesium salt ($\text{Mg}(\text{OCOCH}_2\text{COOC}_2\text{H}_5)_2$) (1.02 g) was added, and the resulting mixture was stirred at room temperature for 18 hours. The reaction mixture was concentrated under reduced pressure, and the residue was dissolved in dichloromethane. The dichloromethane layer was washed with water and dried over magnesium sulfate, and the solvent was evaporated. The residual oil was subjected to column chromatography on silica gel. The fractions eluted with chloroform/methanol (30/1, v/v) gave ethyl 6-(1-imidazolyl)-3-oxohexanoate (0.32 g, 44%) as an oil.

NMR (δ ppm in CDCl_3): 1.28 (3H,t,J=7.4Hz), 2.08 (2H,m), 2.53 (2H,t,J=6.6Hz), 3.41 (2H,s), 4.00 (2H,t,J=6.6Hz), 4.19 (2H,q,J=7.4Hz), 6.91 (1H,s), 7.07 (1H,s), 7.46 (1H,s).

Reference Example 15

[0325] According to the same manner as that described in Reference Example 14, ethyl 6-(1,2,4-triazol-1-yl)-3-oxohexanoate was obtained as an oil from 4-(1,2,4-triazol-1-yl)butyric acid.

NMR (δ ppm in CDCl_3): 1.28 (3H,t,J=7.2Hz), 2.19 (2H,m), 2.59 (2H,t,J=6.6Hz), 3.43 (2H,s), 4.19 (2H,q,J=7.2Hz), 4.23 (2H,t,J=6.6Hz), 7.94 (1H,s), 8.07 (1H,s).


Reference Example 16

[0326] According to the same manner as that described in Reference Example 14, ethyl 7-(1-imidazolyl)-3-oxoheptanoate was obtained as an oil from 5-(1-imidazolyl)valeric acid.

NMR (δ ppm in CDCl_3): 1.27 (3H,t,J=7.4Hz), 1.50-1.90 (4H,m), 2.58 (2H,t,J=6.6Hz), 3.41 (2H,s), 3.95 (2H,t,J=7.0Hz), 4.19 (2H,q,J=7.4Hz), 6.90 (1H,s), 7.06 (1H,s), 7.47 (1H,s).

Claims

1. Use of a compound of the formula (I):



wherein

Y is a nitrogen atom or C-G in which G is an optionally esterified or optionally amidated carboxyl group, or hydroxymethyl group;

R is an optionally substituted hydrocarbon group or optionally substituted heterocyclic group;

X is an oxygen atom or optionally oxidized sulfur atom;

n is 0 or 1;

k is 0 or 1;

G and R may be linked together to form a ring;

each of the ring A and ring B may optionally be substituted;

or a pharmaceutically acceptable salt thereof for formulation of a pharmaceutical composition for inhibiting bone resorption.

2. The use of claim 1, wherein n is 0.

3. The use of claim 1, wherein n is 0 and the hydrocarbon group represented by R is a group of the formula:



wherein X^1 is an oxygen atom, optionally oxidized sulfur atom or $-(\text{CH}_2)_m-$ in which m is an integer of 0 to 5; and Z^1 is an optionally substituted hydrocarbon group, optionally substituted heterocyclic group or optionally substituted amino group.

10

4. The use of claim 3, wherein X^1 is a thio group, sulfinyl group or sulfonyl group.

5. The use of claim 4, wherein X^1 is a thio group.

15 6. The use of claim 3, wherein X^1 is $-(\text{CH}_2)_m-$ and m is 0.

7. The use of claim 3, wherein X^1 is $-(\text{CH}_2)_m-$ and m is 1 or 2.

8. The use of claim 3, wherein the optionally substituted heterocyclic group represented by Z^1 is an aromatic 5-membered heterocyclic group containing 2 to 3 hetero atoms.

20

9. The use of claim 3, wherein Z^1 is an optionally substituted amino group.

10. The use of claim 1, wherein the optionally substituted heterocyclic group represented by R is an optionally substituted 5- to 7-membered heterocyclic group containing one sulfur atom, nitrogen atom or oxygen atom, an optionally substituted 5- to 6-membered heterocyclic group containing 2 to 4 nitrogen atoms, or an optionally substituted 5- to 6-membered heterocyclic group containing 1 to 2 nitrogen atoms and one sulfur atom or oxygen atom.

25

11. The use of claim 1, wherein Y is C-G.

30

12. The use of claim 11, wherein G is a $(\text{C}_{1-6}$ alkyl)oxycarbonyl group.

13. The use of claim 11, wherein G is an ethoxycarbonyl group.

35 14. The use of claim 1, wherein the ring A is substituted with at least one alkoxy group.

15. The use of claim 1, wherein the ring A is substituted with at least one methoxy group.

16. The use of claim 1, wherein the ring A is substituted with the same or different two alkoxy groups.

40

17. The use of claim 1, wherein the ring A is substituted with two methoxy groups.

18. The use of claim 1, wherein the ring A is substituted with two methoxy groups at the 6- and 7-positions of the quinoline ring or quinazoline ring.

45

19. The use of claim 1, wherein the ring B is substituted with at least one alkoxy group.

20. The use of claim 1, wherein the ring B is substituted with at least one methoxy group.

50 21. The use of claim 1, wherein the ring B is substituted with at least one isopropoxy group.

22. The use of claim 1, wherein the ring B is substituted with the same or different two alkoxy groups.

23. The use of claim 1, wherein the ring B is substituted with one methoxy group.

55


24. The use of claim 1, wherein the ring B is substituted with an isopropoxy group at the 3-position and a methoxy group at the 4-position.

25. The use of claim 1, wherein k is 0.

26. The use of claim 1, wherein the compound of the formula (I) is

methyl 4-(3,4-dimethoxyphenyl)-2-ethyl-6,7-dimethoxyquinoline-3-carboxylate;
ethyl 6-chloro-2-methyl-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate;
6,7-dimethoxy-9-phenylfuro[3,4-b]quinoline-1(3H)-one;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate;
4-(3,4-dimethoxyphenyl)-2-(2-hydroxyethylthiomethyl)-6,7-dimethoxyquinazoline;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[(4-methyl-1,2,4-triazol-3-yl)thiomethyl]quinazoline;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[2-(1-methylimidazol-2-yl)ethyl]quinoline-3-carboxylate;
methyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-3-methoxycarbonylquinoline-2-acetate;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
ethyl 4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
ethyl 4-(4-hydroxy-3-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
ethyl 7-hydroxy-6-methoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate 1-oxide; or
ethyl 2-(N,N-diethylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate.

27. Use of a compound of the formula (I):




wherein

Y is a nitrogen atom or C-G in which G is an optionally esterified or optionally amidated carboxyl group, or hydroxymethyl group;
R is an optionally substituted hydrocarbon group or optionally substituted heterocyclic group;
X is an oxygen atom or optionally oxidized sulfur atom;
n is 0 or 1;
k is 0 or 1;
G and R may be linked together to form a ring;
each of the ring A and ring B may optionally be substituted;
or a pharmaceutically acceptable salt thereof for formulation of a pharmaceutical composition for preventing or treating osteoporosis.

28. The use of claim 27, wherein n is 0.

29. The use of claim 27, wherein n is 0 and the hydrocarbon group represented by R is a group of the formula:




wherein X^1 is an oxygen atom, optionally oxidized sulfur atom or $-(\text{CH}_2)_m-$ in which m is an integer of 0 to 5; and Z^1 is an optionally substituted hydrocarbon group, optionally substituted heterocyclic group or optionally substituted amino group.

30. The use of claim 29, wherein X^1 is a thio group, sulfinyl group or sulfonyl group.

31. The use of claim 30, wherein X¹ is a thio group.
32. The use of claim 29, wherein X¹ is -(CH₂)_m- and m is 0.
- 5 33. The use of claim 29, wherein X¹ is -(CH₂)_m- and m is 1 or 2.
34. The use of claim 29, wherein the optionally substituted heterocyclic group represented by Z¹ is an aromatic 5-membered heterocyclic group containing 2 to 3 hetero atoms.
- 10 35. The use of claim 29, wherein Z¹ is an optionally substituted amino group.
36. The use of claim 27, wherein the optionally substituted heterocyclic group represented by R is an optionally substituted 5- to 7-membered heterocyclic group containing one sulfur atom, nitrogen atom or oxygen atom, an optionally substituted 5- to 6-membered heterocyclic group containing 2 to 4 nitrogen atoms, or an optionally substituted 5-
15 to 6-membered heterocyclic group containing 1 to 2 nitrogen atoms and one sulfur atom or oxygen atom.
37. The use of claim 27, wherein Y is C-G.
38. The use of claim 37, wherein G is a (C₁₋₆ alkyl)oxycarbonyl group.
- 20 39. The use of claim 37, wherein G is an ethoxycarbonyl group.
40. The use of claim 27, wherein the ring A is substituted with at least one alkoxy group.
- 25 41. The use of claim 27, wherein the ring A is substituted with at least one methoxy group.
42. The use of claim 27, wherein the ring A is substituted with the same or different two alkoxy groups.
43. The use of claim 27, wherein the ring A is substituted with two methoxy groups.
- 30 44. The use of claim 27, wherein the ring A is substituted with two methoxy groups at the 6- and 7-positions of the quinoline ring or quinazoline ring.
45. The use of claim 27, wherein the ring B is substituted with at least one alkoxy group.
- 35 46. The use of claim 27, wherein the ring B is substituted with at least one methoxy group.
47. The use of claim 27, wherein the ring B is substituted with at least one isopropoxy group.
- 40 48. The use of claim 27, wherein the ring B is substituted with the same or different two alkoxy groups.
49. The use of claim 27, wherein the ring B is substituted with one methoxy group.
50. The use of claim 27, wherein the ring B is substituted with an isopropoxy group at the 3-position and a methoxy
45 group at the 4-position.
51. The use of claim 27, wherein k is 0.
52. The use of claim 27, wherein the compound of the formula (I) is
50 methyl 4-(3,4-dimethoxyphenyl)-2-ethyl-6,7-dimethoxyquinoline-3-carboxylate;
ethyl 6-chloro-2-methyl-4-(3,4-dimethoxyphenyl)quinoline-3-carboxylate;
6,7-dimethoxy-9-phenylfuro[3,4-b]quinoline-1(3H)-one;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[(1-methylimidazol-2-yl)thiomethyl]quinoline-3-carboxylate;
55 4-(3,4-dimethoxyphenyl)-2-(2-hydroxyethylthiomethyl)-6,7-dimethoxyquinazoline;
4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[(4-methyl-1,2,4-triazol-3-yl)thiomethyl]quinazoline;
ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-[2-(1-methylimidazol-2-yl)ethyl]quinoline-3-carboxylate;
methyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-3-methoxycarbonylquinoline-2-acetate;

ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
 ethyl 4-(3-isopropoxy-4-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
 ethyl 4-(4-hydroxy-3-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
 ethyl 7-hydroxy-6-methoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate;
 ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate 1-oxide; or
 ethyl 2-(N,N-diethylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxyquinoline-3-carboxylate.

53. A compound of the formula (I'):




wherein

Y is C-G in which G is an optionally amidated carboxyl group, or hydroxymethyl group;
 R is an optionally substituted hydrocarbon-group or optionally substituted heterocyclic group;
 X is an oxygen atom or optionally oxidized sulfur atom;
 n is 0 or 1;
 k is 0 or 1;
 G and R may be linked together to form a ring;
 each of the ring A and B may optionally be substituted;
 or a pharmaceutically acceptable salt thereof.

54. The compound of claim 53, wherein n is 0.

55. The compound of claim 53, wherein n is 0 and the hydrocarbon group represented by R is a group of the formula:



wherein X^1 is an oxygen atom, optionally oxidized sulfur atom or $-(\text{CH}_2)_m-$ in which m is an integer of 0 to 5; and Z^1 is an optionally substituted hydrocarbon group, optionally substituted heterocyclic group or optionally substituted amino group.

56. The compound of claim 55, wherein X^1 is a thio group, sulfinyl group or sulfonyl group.

57. The compound of claim 56, wherein X^1 is a thio group.

58. The compound of claim 55, wherein X^1 is $-(\text{CH}_2)_m-$ and m is 0.

59. The compound of claim 55, wherein X^1 is $-(\text{CH}_2)_m-$ and m is 1 or 2.



60. The compound of claim 55, wherein the optionally substituted heterocyclic group represented by Z^1 is an aromatic 5-membered heterocyclic group containing 2 to 3 hetero atoms.

61. The compound of claim 55, wherein Z^1 is an optionally substituted amino group.

62. The compound of claim 53, wherein the optionally substituted heterocyclic group represented by R is an optionally substituted 5- to 7-membered heterocyclic group containing one sulfur atom, nitrogen atom or oxygen atom, an optionally substituted 5- to 6-membered heterocyclic group containing 2 to 4 nitrogen atoms, or an optionally sub-

stituted 5- to 6-membered heterocyclic group containing 1 to 2 nitrogen atoms and one sulfur atom or oxygen atom.

63. The compound of claim 53, wherein the ring A is substituted with at least one alkoxy group.
- 5 64. The compound of claim 53, wherein the ring A is substituted with at least one methoxy group.
65. The compound of claim 53, wherein the ring A is substituted with the same or different two alkoxy groups.
66. The compound of claim 53, wherein the ring A is substituted with two methoxy groups.
- 10 67. The compound of claim 53, wherein the ring A is substituted with two methoxy groups at the 6- and 7-positions of the quinoline ring or quinazoline ring.
68. The compound of claim 53, wherein the ring B is substituted with at least one alkoxy group.
- 15 69. The compound of claim 53, wherein the ring B is substituted with at least one methoxy group.
70. The compound of claim 53, wherein the ring B is substituted with at least one isopropoxy group.
- 20 71. The compound of claim 53, wherein the ring B is substituted with the same or different two alkoxy groups.
72. The compound of claim 53, wherein the ring B is substituted with one methoxy group.
- 25 73. The compound of claim 53, wherein the ring B is substituted with an isopropoxy group at the 3-position and a methoxy group at the 4-position.
74. The compound of claim 53, wherein k is 0.
- 30 75. The compound of claim 53 having the 5-membered ring formed by G and R, which is represented by the formula of (II) or (III):



45 wherein

R^3 is hydrogen, an optionally substituted hydrocarbon group or optionally substituted heterocyclic group;
k is 0 or 1;

50 Z^1 is an optionally substituted hydrocarbon group, optionally substituted heterocyclic group or optionally substituted amino group;


each of the ring A and ring B may optionally be substituted.

- 55 76. A pharmaceutical composition which comprises a compound of the formula (I') as defined in claim 53 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.
77. A pharmaceutical composition of claim 76, which is for inhibiting bone resorption.

78. A pharmaceutical composition of claim 76, which is for preventing or treating osteoporosis.

Patentansprüche

1. Verwendung einer Verbindung der Formel (I):




worin

Y ein Stickstoffatom oder C-G ist, worin G eine gegebenenfalls veresterte oder gegebenenfalls amidierter Carboxylgruppe oder Hydroxymethylgruppe ist,
 R eine gegebenenfalls substituierte Kohlenwasserstoffgruppe oder gegebenenfalls substituierte heterocyclische Gruppe ist,
 X ein Sauerstoffatom oder gegebenenfalls oxidiertes Schwefelatom ist,
 n 0 oder 1 ist,
 k 0 oder 1 ist,
 G und R unter Bilden eines Ringes zusammen verbunden sein können,
 Ring A und Ring B jeweils gegebenenfalls substituiert sein können,
 oder eines pharmazeutisch annehmbaren Salzes derselben zur Formulierung einer pharmazeutischen Zusammensetzung zum Hemmen der Knochenresorption.

2. Verwendung von Anspruch 1, wobei n 0 ist.

3. Verwendung von Anspruch 1, wobei n 0 ist und die durch R dargestellte Kohlenwasserstoffgruppe eine Gruppe der Formel



ist, worin X^1 ein Sauerstoffatom, gegebenenfalls oxidiertes Schwefelatom oder $-(\text{CH}_2)_m-$ ist, worin m eine ganze Zahl von 0 bis 5 ist, und Z^1 eine gegebenenfalls substituierte Kohlenwasserstoffgruppe, gegebenenfalls substituierte heterocyclische Gruppe oder gegebenenfalls substituierte Aminogruppe ist.

4. Verwendung von Anspruch 3, wobei X^1 eine Thiogruppe, Sulfinylgruppe oder Sulfonylgruppe ist.

5. Verwendung von Anspruch 4, wobei X^1 eine Thiogruppe ist.

6. Verwendung von Anspruch 3, wobei X^1 $-(\text{CH}_2)_m-$ ist und m 0 ist.

7. Verwendung von Anspruch 3, wobei X^1 $-(\text{CH}_2)_m-$ ist und m 1 oder 2 ist.

8. Verwendung von Anspruch 3, wobei die durch Z^1 dargestellte, gegebenenfalls substituierte heterocyclische Gruppe eine 2 bis 3 Heteroatome enthaltende, aromatische, 5-gliedrige heterocyclische Gruppe ist.

9. Verwendung von Anspruch 3, wobei Z^1 eine gegebenenfalls substituierte Aminogruppe ist.

10. Verwendung von Anspruch 1, wobei die durch R dargestellte, gegebenenfalls substituierte heterocyclische Gruppe eine ein Schwefelatom, Stickstoffatom oder Sauerstoffatom enthaltende, gegebenenfalls substituierte, 5- bis 7-

gliedrige heterocyclische Gruppe, eine 2 bis 4 Stickstoffatome enthaltende, gegebenenfalls substituierte, 5- bis 6-gliedrige heterocyclische Gruppe oder eine 1 bis 2 Stickstoffatome und ein Schwefelatom oder Sauerstoffatom enthaltende, gegebenenfalls substituierte, 5- bis 6-gliedrige heterocyclische Gruppe ist.

- 5 11. Verwendung von Anspruch 1, wobei Y C-G ist.
12. Verwendung von Anspruch 11, wobei G eine (C₁₋₆-Alkyl)oxycarbonylgruppe ist.
13. Verwendung von Anspruch 11, wobei G eine Ethoxycarbonylgruppe ist.
- 10 14. Verwendung von Anspruch 1, wobei Ring A mit wenigstens einer Alkoxygruppe substituiert ist.
- 15 15. Verwendung von Anspruch 1, wobei Ring A mit wenigstens einer Methoxygruppe substituiert ist.
- 16 16. Verwendung von Anspruch 1, wobei Ring A mit zwei gleichen oder verschiedenen Alkoxygruppen substituiert ist.
17. Verwendung von Anspruch 1, wobei Ring A mit zwei Methoxygruppen substituiert ist.
- 18 18. Verwendung von Anspruch 1, wobei Ring A mit zwei Methoxygruppen in 6- und 7-Stellung des Chinolinrings oder Chinazolinrings substituiert ist.
19. Verwendung von Anspruch 1, wobei Ring B mit wenigstens einer Alkoxygruppe substituiert ist.
- 20 20. Verwendung von Anspruch 1, wobei Ring B mit wenigstens einer Methoxygruppe substituiert ist.
- 25 21. Verwendung von Anspruch 1, wobei Ring B mit wenigstens einer Isopropoxygruppe substituiert ist.
22. Verwendung von Anspruch 1, wobei Ring B mit zwei gleichen oder verschiedenen Alkoxygruppen substituiert ist.
- 30 23. Verwendung von Anspruch 1, wobei Ring B mit einer Methoxygruppe substituiert ist.
24. Verwendung von Anspruch 1, wobei Ring B in 3-Stellung mit einer Isopropoxygruppe und in 4-Stellung mit einer Methoxygruppe substituiert ist.
- 35 25. Verwendung von Anspruch 1, wobei k 0 ist.
26. Verwendung von Anspruch 1, wobei die Verbindung der Formel (I)
40 4-(3,4-Dimethoxyphenyl)-2-ethyl-6,7-dimethoxychinolin-3-carbonsäuremethylester,
6-Chlor-2-methyl-4-(3,4-dimethoxyphenyl)chinolin-3-carbonsäureethylester,
6,7-Dimethoxy-9-phenylfuro[3,4-b]chinolin-1(3H)-on,
45 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-[(1-methylimidazol-2-yl)thiomethyl]chinolin-3-carbonsäureethylester,
4-(3,4-Dimethoxyphenyl)-2-(2-hydroxyethylthiomethyl)-6,7-dimethoxychinazolin,
50 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-[(4-methyl-1,2,4-triazol-3-yl)thiomethyl]chinazolin,
4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-[2-(1-methylimidazol-2-yl)ethyl]chinolin-3-carbonsäureethylester,
4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-3-methoxycarbonylchinolin-2-essigsäuremethylester,
55 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester,
4-(3-Isopropoxy-4-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethyle-

ster,


4-(4-Hydroxy-3-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester,

7-Hydroxy-6-methoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester,

4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester-1-oxid
oder

2-(N,N-Diethylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxychinolin-3-carbonsäureethylester ist.

27. Verwendung einer Verbindung der Formel (I):



worin

Y ein Stickstoffatom oder C-G ist, worin G eine gegebenenfalls veresterte oder gegebenenfalls amidierter Carboxylgruppe oder Hydroxymethylgruppe ist,

R eine gegebenenfalls substituierte Kohlenwasserstoffgruppe oder gegebenenfalls substituierte heterocyclische Gruppe ist,

X ein Sauerstoffatom oder gegebenenfalls oxidiertes Schwefelatom ist;

n 0 oder 1 ist,

k 0 oder 1 ist,


G und R unter Bilden eines Ringes zusammen verbunden sein können,

Ring A und Ring B jeweils gegebenenfalls substituiert sein können,

oder eines pharmazeutisch annehmbaren Salzes derselben zur Formulierung einer pharmazeutischen Zusammensetzung zum Verhindern oder Behandeln von Osteoporose.

28. Verwendung von Anspruch 27, wobei n 0 ist.

29. Verwendung von Anspruch 27, wobei n 0 ist und die durch R dargestellte Kohlenwasserstoffgruppe eine Gruppe der Formel



ist, worin X^1 ein Sauerstoffatom, gegebenenfalls oxidiertes Schwefelatom oder $-(\text{CH}_2)_m-$ ist, worin m eine ganze Zahl von 0 bis 5 ist, und Z^1 eine gegebenenfalls substituierte Kohlenwasserstoffgruppe, gegebenenfalls substituierte heterocyclische Gruppe oder gegebenenfalls substituierte Aminogruppe ist.

30. Verwendung von Anspruch 29, wobei X^1 eine Thiogruppe, Sulfinylgruppe oder Sulfonylgruppe ist.

31. Verwendung von Anspruch 30, wobei X^1 eine Thiogruppe ist.


32. Verwendung von Anspruch 29, wobei X^1 $-(\text{CH}_2)_m-$ ist und m 0 ist.

33. Verwendung von Anspruch 29, wobei X^1 $-(\text{CH}_2)_m-$ ist und m 1 oder 2 ist.

34. Verwendung von Anspruch 29, wobei die durch Z^1 dargestellte, gegebenenfalls substituierte heterocyclische Gruppe eine 2 bis 3 Heteroatome enthaltende, aromatische, 5-gliedrige heterocyclische Gruppe ist.
35. Verwendung von Anspruch 29, wobei Z^1 eine gegebenenfalls substituierte Aminogruppe ist.
36. Verwendung von Anspruch 27, wobei die durch R dargestellte, gegebenenfalls substituierte heterocyclische Gruppe eine ein Schwefelatom, Stickstoffatom oder Sauerstoffatom enthaltende, gegebenenfalls substituierte, 5- bis 7-gliedrige heterocyclische Gruppe, eine 2 bis 4 Stickstoffatome enthaltende, gegebenenfalls substituierte, 5- bis 6-gliedrige heterocyclische Gruppe oder eine 1 bis 2 Stickstoffatome und ein Schwefelatom oder Sauerstoffatom enthaltende, gegebenenfalls substituierte, 5- bis 6-gliedrige heterocyclische Gruppe ist.
37. Verwendung von Anspruch 27, wobei Y C-G ist.
38. Verwendung von Anspruch 37, wobei G eine (C_{1-6} -Alkyl)oxycarbonylgruppe ist.
39. Verwendung von Anspruch 37, wobei G eine Ethoxycarbonylgruppe ist.
40. Verwendung von Anspruch 27, wobei Ring A mit wenigstens einer Alkoxygruppe substituiert ist.
41. Verwendung von Anspruch 27, wobei Ring A mit wenigstens einer Methoxygruppe substituiert ist.
42. Verwendung von Anspruch 27, wobei Ring A mit zwei gleichen oder verschiedenen Alkoxygruppen substituiert ist.
43. Verwendung von Anspruch 27, wobei Ring A mit zwei Methoxygruppen substituiert ist.
44. Verwendung von Anspruch 27, wobei Ring A mit zwei Methoxygruppen in 6- und 7-Stellung des Chinolinrings oder Chinazolinrings substituiert ist.
45. Verwendung von Anspruch 27, wobei Ring B mit wenigstens einer Alkoxygruppe substituiert ist.
46. Verwendung von Anspruch 27, wobei Ring B mit wenigstens einer Methoxygruppe substituiert ist.
47. Verwendung von Anspruch 27, wobei Ring B mit wenigstens einer Isopropoxygruppe substituiert ist.
48. Verwendung von Anspruch 27, wobei Ring B mit zwei gleichen oder verschiedenen Alkoxygruppen substituiert ist.
49. Verwendung von Anspruch 27, wobei Ring B mit einer Methoxygruppe substituiert ist.
50. Verwendung von Anspruch 27, wobei Ring B in 3-Stellung mit einer Isopropoxygruppe und in 4-Stellung mit einer Methoxygruppe substituiert ist.
51. Verwendung von Anspruch 27, wobei k 0 ist.
52. Verwendung von Anspruch 27, wobei die Verbindung der Formel (I)
- 4-(3,4-Dimethoxyphenyl)-2-ethyl-6,7-dimethoxychinolin-3-carbonsäuremethylester,
- 6-Chlor-2-methyl-4-(3,4-dimethoxyphenyl)chinolin-3-carbonsäureethylester,
- 6,7-Dimethoxy-9-phenylfuro[3,4-b]chinolin-1(3H)-on,
- 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-[(1-methylimidazol-2-yl)thiomethyl]chinolin-3-carbonsäureethylester,
- 4-(3,4-Dimethoxyphenyl)-2-(2-hydroxyethylthiomethyl)-6,7-dimethoxychinazolin,
- 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-[(4-methyl-1,2,4-triazol-3-yl)thiomethyl]chinazolin,

4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-[2-(1-methylimidazol-2-yl)ethyl]chinolin-3-carbonsäureethylester,
 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-3-methoxycarbonylchinolin-2-essigsäuremethylester,
 5 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester,
 4-(3-Isopropoxy-4-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester,
 10 4-(4-Hydroxy-3-methoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester,
 7-Hydroxy-6-methoxy-4-(3,4-dimethoxyphenyl)-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester,
 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)chinolin-3-carbonsäureethylester-1-oxid
 15 oder
 2-(N,N-Diethylaminomethyl)-4-(3,4-dimethoxyphenyl)-6,7-dimethoxychinolin-3-carbonsäureethylester ist.

53. Verbindung der Formel (I'):




worin

Y C-G ist, worin G eine gegebenenfalls amidierter Carboxylgruppe oder Hydroxymethylgruppe ist,
 R eine gegebenenfalls substituierte Kohlenwasserstoffgruppe oder gegebenenfalls substituierte heterocyclische Gruppe ist,
 X ein Sauerstoffatom oder gegebenenfalls oxidiertes Schwefelatom ist,
 n 0 oder 1 ist,
 k 0 oder 1 ist,
 G und R unter Bilden eines Ringes zusammen verbunden sein können,
 Ring A und Ring B jeweils gegebenenfalls substituiert sein können,
 oder ein pharmazeutisch annehmbares Salz derselben.

54. Verbindung von Anspruch 53, wobei n 0 ist.

55. Verbindung von Anspruch 53, wobei n 0 ist und die durch R dargestellte Kohlenwasserstoffgruppe eine Gruppe der Formel





ist, worin X^1 ein Sauerstoffatom, gegebenenfalls oxidiertes Schwefelatom oder $-(\text{CH}_2)_m-$ ist, worin m eine ganze Zahl von 0 bis 5 ist und Z^1 eine gegebenenfalls substituierte Kohlenwasserstoffgruppe, gegebenenfalls substituierte heterocyclische Gruppe oder gegebenenfalls substituierte Aminogruppe ist.

56. Verbindung von Anspruch 55, wobei X^1 eine Thiogruppe, Sulfinylgruppe oder Sulfonylgruppe ist.

57. Verwendung von Anspruch 56, wobei X^1 eine Thiogruppe ist.

58. Verbindung von Anspruch 55, wobei X^1 $-(CH_2)_m$ - ist und m 0 ist.
59. Verbindung von Anspruch 55, wobei X^1 $-(CH_2)_m$ - ist und m 1 oder 2 ist.
- 5 60. Verbindung von Anspruch 55, wobei die durch Z^1 dargestellte, gegebenenfalls substituierte heterocyclische Gruppe eine 2 bis 3 Heteroatome enthaltende, aromatische, 5-gliedrige heterocyclische Gruppe ist.
61. Verbindung von Anspruch 55, wobei Z^1 eine gegebenenfalls substituierte Aminogruppe ist.
- 10 62. Verbindung von Anspruch 53, wobei die durch R dargestellte, gegebenenfalls substituierte heterocyclische Gruppe eine ein Schwefelatom, Stickstoffatom oder Sauerstoffatom enthaltende, gegebenenfalls substituierte, 5- bis 7-gliedrige heterocyclische Gruppe, eine 2 bis 4 Stickstoffatome enthaltende, gegebenenfalls substituierte, 5- bis 6-gliedrige heterocyclische Gruppe oder eine 1 bis 2 Stickstoffatome und ein Schwefelatom oder Sauerstoffatom enthaltende, gegebenenfalls substituierte, 5- bis 6-gliedrige heterocyclische Gruppe ist.
- 15 63. Verbindung von Anspruch 53, wobei Ring A mit wenigstens einer Alkoxygruppe substituiert ist.
64. Verbindung von Anspruch 53, wobei Ring A mit wenigstens einer Methoxygruppe substituiert ist.
- 20 65. Verbindung von Anspruch 53, wobei Ring A mit zwei gleichen oder verschiedenen Alkoxygruppen substituiert ist.
66. Verbindung von Anspruch 53, wobei Ring A mit zwei Methoxygruppen substituiert ist.
- 25 67. Verbindung von Anspruch 53, wobei Ring A mit zwei Methoxygruppen in 6- und 7-Stellung des Chinolinrings oder Chinazolinrings substituiert ist.
68. Verbindung von Anspruch 53, wobei Ring B mit wenigstens einer Alkoxygruppe substituiert ist.
69. Verbindung von Anspruch 53, wobei Ring B mit wenigstens einer Methoxygruppe substituiert ist.
- 30 70. Verbindung von Anspruch 53, wobei Ring B mit wenigstens einer Isopropoxygruppe substituiert ist.
71. Verbindung von Anspruch 53, wobei Ring B mit zwei gleichen oder verschiedenen Alkoxygruppen substituiert ist.
- 35 72. Verbindung von Anspruch 53, wobei Ring B mit einer Methoxygruppe substituiert ist.
73. Verbindung von Anspruch 53, wobei Ring B in 3-Stellung mit einer Isopropoxygruppe und in 4-Stellung mit einer Methoxygruppe substituiert ist.
- 40 74. Verbindung von Anspruch 53, wobei k 0 ist.
75. Verbindung von Anspruch 53 mit einem durch G und R gebildeten 5-gliedrigen Ring, die durch die Formel (II) oder (III) dargestellt wird:



worin

R³ Wasserstoff, eine gegebenenfalls substituierte Kohlenwasserstoffgruppe oder gegebenenfalls substituierte heterocyclische Gruppe ist,

k 0 oder 1 ist,

Z¹ eine gegebenenfalls substituierte Kohlenwasserstoffgruppe, gegebenenfalls substituierte heterocyclische Gruppe oder gegebenenfalls substituierte Aminogruppe ist,

Ring A und Ring B jeweils gegebenenfalls substituiert sein können.


76. Pharmazeutische Zusammensetzung, die eine in Anspruch 53 definierte Verbindung der Formel (I') oder ein pharmazeutisch annehmbares Salz derselben und einen pharmazeutisch annehmbaren Träger umfaßt.

- 77. Pharmazeutische Zusammensetzung von Anspruch 76 zum Hemmen der Knochenresorption.**

- 78. Pharmazeutische Zusammensetzung von Anspruch 76 zum Verhindern oder Behandeln von Osteoporose.**

Revendications

1. Emploi d'un composé de formule (I) :



dans laquelle

Y représente un atome d'azote, ou C-G où G représente un groupe carboxyle, éventuellement estérifié ou amidifié, ou un groupe hydroxyméthyle ;

R représente un groupe hydrocarboné, portant éventuellement un substituant, ou un groupe hétérocyclique, portant éventuellement un substituant ;

X représente un atome d'oxygène, ou un atome de soufre, éventuellement oxydé ;

n vaut 0 ou 1 :

k vaut 0 ou 1 :


G et R peuvent être raccordés pour former un cycle ;

et chacun des cycles A et B peut éventuellement porter un substituant :

ou d'un sel d'un tel composé, admissible en pharmacie, dans la formulation d'une composition pharmaceutique destinée à empêcher la résorption des os.

2. Emploi conforme à la revendication 1, où n vaut 0.

3. Emploi conforme à la revendication 1, où n vaut 0 et le groupe hydrocarboné représenté par R est un groupe de formule




dans laquelle X^1 représente un atome d'oxygène, un atome de soufre éventuellement oxydé, ou un groupe $-(CH_2)_m-$ où m vaut de 0 à 5, et Z^1 représente un groupe hydrocarboné portant éventuellement un substituant, un groupe hétérocyclique portant éventuellement un substituant, ou un groupe amino portant éventuellement un substituant.

4. Emploi conforme à la revendication 3, où X^1 représente un groupe thio, sulfinyle ou sulfonyle.
5. Emploi conforme à la revendication 4, où X^1 représente un groupe thio.
- 5 6. Emploi conforme à la revendication 3, où X^1 représente un groupe $-(CH_2)_m-$ et m vaut 0.
7. Emploi conforme à la revendication 3, où X^1 représente un groupe $-(CH_2)_m-$ et m vaut 1 ou 2.
8. Emploi conforme à la revendication 3, où le groupe hétérocyclique portant éventuellement un substituant et représenté par Z^1 est un groupe hétérocyclique aromatique à 5 chaînons, qui comporte deux ou trois hétéroatomes.
- 10 9. Emploi conforme à la revendication 3, où Z^1 représente un groupe amino portant éventuellement un substituant.
- 15 10. Emploi conforme à la revendication 1, où le groupe hétérocyclique portant éventuellement un substituant et représenté par R est un groupe hétérocyclique à 5-7 chaînons, qui porte éventuellement un substituant et qui comporte un seul atome de soufre, d'azote ou d'oxygène, un groupe hétérocyclique à 5-6 chaînons, qui porte éventuellement un substituant et qui comporte de deux à quatre atomes d'azote, ou un groupe hétérocyclique à 5-6 chaînons, qui porte éventuellement un substituant et qui comporte un ou deux atomes d'azote et un seul atome de soufre ou d'oxygène.
- 20 11. Emploi conforme à la revendication 1, où Y représente C-G.
12. Emploi conforme à la revendication 11, où G représente un groupe (alcoxy en C_{1-6})carbonyle.
- 25 13. Emploi conforme à la revendication 11, où G représente un groupe éthoxycarbonyle.
14. Emploi conforme à la revendication 1, où le cycle A porte au moins un groupe alcoxy comme substituant.
15. Emploi conforme à la revendication 1, où le cycle A porte au moins un groupe méthoxy comme substituant.
- 30 16. Emploi conforme à la revendication 1, où le cycle A porte comme substituants deux groupes alcoxy identiques ou différents.
17. Emploi conforme à la revendication 1, où le cycle A porte deux groupes méthoxy comme substituants.
- 35 18. Emploi conforme à la revendication 1, où le cycle A porte deux groupes méthoxy comme substituants, en positions 6 et 7 du cycle quinoline ou quinazoline.
19. Emploi conforme à la revendication 1, où le cycle B porte au moins un groupe alcoxy comme substituant.
- 40 20. Emploi conforme à la revendication 1, où le cycle B porte au moins un groupe méthoxy comme substituant.
21. Emploi conforme à la revendication 1, où le cycle B porte au moins un groupe isopropoxy comme substituant.
- 45 22. Emploi conforme à la revendication 1, où le cycle B porte comme substituants deux groupes alcoxy identiques ou différents.
23. Emploi conforme à la revendication 1, où le cycle B porte un seul groupe méthoxy comme substituant.
- 50 24. Emploi conforme à la revendication 1, où le cycle B porte comme substituants un groupe isopropoxy en position 3 et un groupe méthoxy en position 4.
25. Emploi conforme à la revendication 1, où k vaut 0.
- 55 26. Emploi conforme à la revendication 1, où le composé de formule (I) est l'un des suivants :

4-(3,4-diméthoxyphényl)-2-éthyl-6,7-diméthoxyquinoline-3-carboxylate de méthyle ;
6-chloro-2-méthyl-4-(3,4-diméthoxyphényl)quinoline-3-carboxylate d'éthyle ;

6,7-diméthoxy-9-phényl-furo[3,4-b]quinoline-1(3H)-one ;
 4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-[(1-méthyl-imidazole-2-yl)thiométhyl]quinoline-3-carboxylate
 d'éthyle ;
 4-(3,4-diméthoxyphényl)-2-(2-hydroxyéthylthiométhyl)-6,7-diméthoxyquinazoline ;
 4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-[(4-méthyl-1,2,4-triazole-3-yl)thiométhyl]quinazoline ;
 4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-[2-(1-méthyl-imidazole-2-yl)éthyl]quinoline-3-carboxylate d'éthyle ;
 4-(3,4-diméthoxyphényl)-6,7-diméthoxy-3-méthoxycarbonyl-quinoline-2-acétate de méthyle ;
 4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate d'éthyle ;
 4-(3-isopropoxy-4-méthoxyphényl)-6,7-diméthoxy-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate
 d'éthyle ;
 4-(4-hydroxy-3-méthoxyphényl)-6,7-diméthoxy-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate d'éthyle ;
 7-hydroxy-6-méthoxy-4-(3,4-diméthoxyphényl)-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate d'éthyle ;
 1-oxyde de 4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate
 d'éthyle ; et
 2-(N,N-diéthylaminométhyl)-4-(3,4-diméthoxyphényl)-6,7-diméthoxyquinoline-3-carboxylate d'éthyle.

27. Emploi d'un composé de formule (I) :




dans laquelle

Y représente un atome d'azote, ou C-G où G représente un groupe carboxyle, éventuellement estérifié ou ami-
 difié, ou un groupe hydroxyméthyle ;
 R représente un groupe hydrocarboné, portant éventuellement un substituant, ou un groupe hétérocyclique,
 portant éventuellement un substituant ;
 X représente un atome d'oxygène, ou un atome de soufre, éventuellement oxydé ;
 n vaut 0 ou 1 ;
 k vaut 0 ou 1 ;
 G et R peuvent être raccordés pour former un cycle ;
 et chacun des cycles A et B peut éventuellement porter un substituant ;
 ou d'un sel d'un tel composé, admissible en pharmacie, dans la formulation d'une composition pharmaceuti-
 que destinée à la prévention ou au traitement de l'ostéoporose.

28. Emploi conforme à la revendication 27, où n vaut 0.

29. Emploi conforme à la revendication 27, où n vaut 0 et le groupe hydrocarboné représenté par R est un groupe de
 formule



dans laquelle X^1 représente un atome d'oxygène, un atome de soufre éventuellement oxydé, ou un groupe -
 $(\text{CH}_2)_m$ - où m vaut de 0 à 5, et Z^1 représente un groupe hydrocarboné portant éventuellement un substituant, un
 groupe hétérocyclique portant éventuellement un substituant, ou un groupe amino portant éventuellement un
 substituant.

30. Emploi conforme à la revendication 29, où X^1 représente un groupe thio, sulfinyle ou sulfonyle.

31. Emploi conforme à la revendication 30, où X^1 représente un groupe thio.
32. Emploi conforme à la revendication 29, où X^1 représente un groupe $-(CH_2)_m-$ et m vaut 0.
- 5 33. Emploi conforme à la revendication 29, où X^1 représente un groupe $-(CH_2)_m-$ et m vaut 1 ou 2.
34. Emploi conforme à la revendication 29, où le groupe hétérocyclique portant éventuellement un substituant et représenté par Z^1 est un groupe hétérocyclique aromatique à 5 chaînons, qui comporte deux ou trois hétéroatomes.
- 10 35. Emploi conforme à la revendication 29, où Z^1 représente un groupe amino portant éventuellement un substituant.
36. Emploi conforme à la revendication 27, où le groupe hétérocyclique portant éventuellement un substituant et représenté par R est un groupe hétérocyclique à 5-7 chaînons, qui porte éventuellement un substituant et qui comporte un seul atome de soufre, d'azote ou d'oxygène, un groupe hétérocyclique à 5-6 chaînons, qui porte éventuellement un substituant et qui comporte de deux à quatre atomes d'azote, ou un groupe hétérocyclique à 5-6 chaînons, qui porte éventuellement un substituant et qui comporte un ou deux atomes d'azote et un seul atome de soufre ou d'oxygène.
- 15 37. Emploi conforme à la revendication 27, où Y représente C-G.
- 20 38. Emploi conforme à la revendication 37, où G représente un groupe (alcoxy en C_{1-6})carbonyle.
39. Emploi conforme à la revendication 37, où G représente un groupe éthoxycarbonyle.
- 25 40. Emploi conforme à la revendication 27, où le cycle A porte au moins un groupe alcoxy comme substituant.
41. Emploi conforme à la revendication 27, où le cycle A porte au moins un groupe méthoxy comme substituant.
- 30 42. Emploi conforme à la revendication 27, où le cycle A porte comme substituants deux groupes alcoxy identiques ou différents.
43. Emploi conforme à la revendication 27, où le cycle A porte deux groupes méthoxy comme substituants.
- 35 44. Emploi conforme à la revendication 27, où le cycle A porte deux groupes méthoxy comme substituants, en positions 6 et 7 du cycle quinoline ou quinazoline.
45. Emploi conforme à la revendication 27, où le cycle B porte au moins un groupe alcoxy comme substituant.
46. Emploi conforme à la revendication 27, où le cycle B porte au moins un groupe méthoxy comme substituant.
- 40 47. Emploi conforme à la revendication 27, où le cycle B porte au moins un groupe isopropoxy comme substituant.
48. Emploi conforme à la revendication 27, où le cycle B porte comme substituants deux groupes alcoxy identiques ou différents.
- 45 49. Emploi conforme à la revendication 27, où le cycle B porte un seul groupe méthoxy comme substituant.
50. Emploi conforme à la revendication 27, où le cycle B porte comme substituants un groupe isopropoxy en position 3 et un groupe méthoxy en position 4.
- 50 51. Emploi conforme à la revendication 27, où k vaut 0.
52. Emploi conforme à la revendication 27, où le composé de formule (I) est l'un des suivants :
- 55 4-(3,4-diméthoxyphényl)-2-éthyl-6,7-diméthoxyquinoline-3-carboxylate de méthyle ;
 6-chloro-2-méthyl-4-(3,4-diméthoxyphényl)quinoline-3-carboxylate d'éthyle ;
 6,7-diméthoxy-9-phényl-furo[3,4-b]quinoline-1(3H)-one ;
 4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-[(1-méthyl-imidazole-2-yl)thiométhyl]quinoline-3-carboxylate

d'éthyle ;

4-(3,4-diméthoxyphényl)-2-(2-hydroxyéthylthiométhyl)-6,7-diméthoxyquinazoline ;

4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-[(4-méthyl-1,2,4-triazole-3-yl)thiométhyl]quinazoline ;

4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-[2-(1-méthyl-imidazole-2-yl)éthyl]quinoline-3-carboxylate d'éthyle ;

4-(3,4-diméthoxyphényl)-6,7-diméthoxy-3-méthoxycarbonyl-quinoline-2-acétate de méthyle ;

4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate d'éthyle ;

4-(3-isopropoxy-4-méthoxyphényl)-6,7-diméthoxy-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate d'éthyle ;


4-(4-hydroxy-3-méthoxyphényl)-6,7-diméthoxy-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate d'éthyle ;

7-hydroxy-6-méthoxy-4-(3,4-diméthoxyphényl)-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate d'éthyle ;

1-oxyde de 4-(3,4-diméthoxyphényl)-6,7-diméthoxy-2-(1,2,4-triazole-1-yl-méthyl)quinoline-3-carboxylate d'éthyle ; et

2-(N,N-diéthylaminométhyl)-4-(3,4-diméthoxyphényl)-6,7-diméthoxyquinoline-3-carboxylate d'éthyle.

53. Composé de formule (I') :



dans laquelle

Y représente C-G où G représente un groupe carboxyle, éventuellement amidifié, ou un groupe hydroxyméthyle ;

R représente un groupe hydrocarboné, portant éventuellement un substituant, ou un groupe hétérocyclique, portant éventuellement un substituant ;

X représente un atome d'oxygène, ou un atome de soufre, éventuellement oxydé ;

n vaut 0 ou 1 ;

k vaut 0 ou 1 ;


G et R peuvent être raccordés pour former un cycle ;

et chacun des cycles A et B peut éventuellement porter un substituant ;

ou sel d'un tel composé, admissible en pharmacie.

54. Composé conforme à la revendication 53, où n vaut 0.

55. Composé conforme à la revendication 53, où n vaut 0 et le groupe hydrocarboné représenté par R est un groupe de formule



dans laquelle X^1 représente un atome d'oxygène, un atome de soufre éventuellement oxydé, ou un groupe $-(\text{CH}_2)_m-$ où m vaut de 0 à 5, et Z^1 représente un groupe hydrocarboné portant éventuellement un substituant, un groupe hétérocyclique portant éventuellement un substituant, ou un groupe amino portant éventuellement un substituant.

56. Composé conforme à la revendication 55, où X^1 représente un groupe thio, sulfinyle ou sulfonyle.



57. Composé conforme à la revendication 56, où X^1 représente un groupe thio.

58. Composé conforme à la revendication 55, où X^1 représente un groupe $-(\text{CH}_2)_m-$ et m vaut 0.

59. Composé conforme à la revendication 55, où X^1 représente un groupe $-(CH_2)_m-$ et m vaut 1 ou 2.
- 5 60. Composé conforme à la revendication 55, où le groupe hétérocyclique portant éventuellement un substituant et représenté par Z^1 est un groupe hétérocyclique aromatique à 5 chaînons, qui comporte deux ou trois hétéroatomes.
61. Composé conforme à la revendication 55, où Z^1 représente un groupe amino portant éventuellement un substituant.
- 10 62. Composé conforme à la revendication 53, où le groupe hétérocyclique portant éventuellement un substituant et représenté par R est un groupe hétérocyclique à 5-7 chaînons, qui porte éventuellement un substituant et qui comporte un seul atome de soufre, d'azote ou d'oxygène, un groupe hétérocyclique à 5-6 chaînons, qui porte éventuellement un substituant et qui comporte de deux à quatre atomes d'azote, ou un groupe hétérocyclique à 5-6 chaînons, qui porte éventuellement un substituant et qui comporte un ou deux atomes d'azote et un seul atome de soufre ou d'oxygène.
- 15 63. Composé conforme à la revendication 53, où le cycle A porte au moins un groupe alcoxy comme substituant.
64. Composé conforme à la revendication 53, où le cycle A porte au moins un groupe méthoxy comme substituant.
- 20 65. Composé conforme à la revendication 53, où le cycle A porte comme substituants deux groupes alcoxy identiques ou différents.
66. Composé conforme à la revendication 53, où le cycle A porte deux groupes méthoxy comme substituants.
- 25 67. Composé conforme à la revendication 53, où le cycle A porte deux groupes méthoxy comme substituants, en positions 6 et 7 du cycle quinoline ou quinazoline.
68. Composé conforme à la revendication 53, où le cycle B porte au moins un groupe alcoxy comme substituant.
- 30 69. Composé conforme à la revendication 53, où le cycle B porte au moins un groupe méthoxy comme substituant.
70. Composé conforme à la revendication 53, où le cycle B porte au moins un groupe isopropoxy comme substituant.
- 35 71. Composé conforme à la revendication 53, où le cycle B porte comme substituants deux groupes alcoxy identiques ou différents.
72. Composé conforme à la revendication 53, où le cycle B porte un groupe méthoxy comme substituant.
- 40 73. Composé conforme à la revendication 53, où le cycle B porte comme substituants un groupe isopropoxy en position 3 et un groupe méthoxy en position 4.
74. Composé conforme à la revendication 53, où k vaut 0.
- 45 75. Composé conforme à la revendication 53, qui comporte un cycle à 5 chaînons formé par G et R et qui est représenté par l'une des formules (II) et (III) :

50

55



dans lesquelles

R^3 représente un atome d'hydrogène, un groupe hydrocarboné, portant éventuellement un substituant, ou un groupe hétérocyclique, portant éventuellement un substituant ;

k vaut 0 ou 1 ;

Z^1 représente un groupe hydrocarboné, portant éventuellement un substituant, un groupe hétérocyclique, portant éventuellement un substituant, ou un groupe amino, portant éventuellement un substituant ;
et chacun des cycles A et B peut éventuellement porter un substituant.

76. Composition pharmaceutique comprenant un composé de formule (I'), conforme à la revendication 53, ou un sel d'un tel composé, admissible en pharmacie, et un véhicule admissible en pharmacie.

77. Composition pharmaceutique conforme à la revendication 76, destinée à empêcher la résorption des os.

78. Composition pharmaceutique conforme à la revendication 76, destinée à la prévention ou au traitement de l'ostéoporose.